


# Estrogen receptor beta transcriptional regulation: A potential mechanism for colon cancer prevention



Rajitha Indukuri

# **Estrogen receptor beta transcriptional regulation: A potential mechanism for colon cancer prevention**

Rajitha Indukuri

Karolinska Institutet

Department of Biosciences and Nutrition

KTH Royal Institute of Technology

School of Engineering Sciences in Chemistry, Biotechnology and Health

Department of Protein Sciences



Stockholm 2021

All previously published papers were reproduced with permission from the publisher.

Published by Karolinska Institutet.

Printed by Universitetsservice US-AB

© Rajitha Indukuri, 2021

ISBN 978-91-8016-227-2

# Estrogen receptor beta transcriptional regulation: a potential mechanism for colon cancer prevention

## THESIS FOR DOCTORAL DEGREE (Ph.D.)

By

**Rajitha Indukuri**

The thesis will be defended digitally and in public at Karolinska Institute and Royal Institute of Technology will be held at Karolinska Institute Campus Flemingsberg, Neo, DNA lecture hall, floor 5, Blickagången 16, Huddinge.

Friday, May 28<sup>th</sup>, 2021 at 9.30

*Principal Supervisor:*

Cecilia Williams  
Karolinska Institute  
Department of Biosciences and Nutrition

*Co-supervisor(s):*

Afshin Ahmadian  
Royal Institute of Technology, KTH  
Department of Gene Technology

*Opponent:*

Jason Carroll  
Cambridge Institute  
Department of Oncology

*Examination Board:*

Ann-Kristin Östlund Farrants  
Stockholm University  
Department of Molecular Biosciences

Monica Nistér  
Karolinska Institute  
Department of Oncology and Pathology

Eckardt Treuter  
Karolinska Institute  
Department of Biosciences and Nutrition



To my parents and  
my father-in-law



## ABSTRACT

Colorectal cancer (CRC) is the third leading cause of death from cancer in both men and women in the Western world. Improved screening efforts, surveillance, and treatment have reduced CRC mortality in older patients. However, the incidence is increasing in young adults, even in the absence of CRC family history. This may indicate an influence of increasing obesity, changed dietary patterns, and lifestyle factors. The progression of CRC is a multistep procedure that takes 10-15 years, thus offering a time to implement preventative measures and early detection. There is a critical need to develop more effective preventive therapies due to the risks posed by current prevention therapies. The best CRC prophylactic agent should be both safe and suitable to use for a long time (1).

In preclinical studies, estrogen has been shown to protect from CRC, and substantial evidence suggests it is through estrogen receptor beta (ER $\beta$ ). Natural ER $\beta$  selective agonists have been tested in phase II clinical trials to treat menopause symptoms and proven to be safe and well-tolerated with no side effects (2, 3). Thus, selective activation of ER $\beta$  with selective agonists, which do not activate estrogen receptor alpha (ER $\alpha$ ), is a potential clinical approach in preventing adenomatous polyps progression into CRC. However, the mechanism of these beneficial ER $\beta$  effects is not well understood, and there is a significant knowledge gap in this area.

The overall aim of this thesis was to identify the mechanistic background of the intestinal ER $\beta$ -mediated antitumorigenic effects in the colon and further to explore ER $\beta$  as a preventative approach in CRC. One specific aim was to determine whether ER $\beta$  present specifically in colon epithelium is responsible for protecting from CRC, which is addressed in Paper I. To understand the impact of ER $\beta$  in protecting from colitis-associated CRC (CA-CRC), we have induced colitis in intestinal-specific ER $\beta$  knockout mice of both sexes. The loss of intestinal ER $\beta$  aggravated CA-CRC in a sex-dependent manner. The incidence of tumors increased in males, while in females, the size of the tumors was enhanced. We identified that ER $\beta$  attenuates tumor necrosis factor alpha (TNF $\alpha$ ) induced epithelial cell damage and modulates the regulation of key nuclear factor- $\kappa$ B (NF $\kappa$ B) members. As a direct consequence, ER $\beta$  was found to reduce inflammation and to control intestinal crypt cell proliferation.

Another aim was to explore transcriptional regulation by ER $\beta$  through mapping of chromatin binding sites and interaction with NF $\kappa$ B, which is studied in Paper II and IV. Commonly used ER $\beta$  antibodies have been shown to be unspecific towards ER $\beta$ ; this study used a validated ER $\beta$  antibody to map genome-wide ER $\beta$  binding sites in colon cancer cells. We observed that the presence of ER $\beta$  also modulated the regulatory chromatin mark H3K27AC in potential enhancers of transcriptional regulation, Wnt signaling, and cell proliferation. Further, motif analysis indicated a novel ER $\beta$  colon-specific cross-talk with TCF, and KLF motifs supported a interaction between  $\beta$ -catenin/TCF and ER $\beta$ . We found that ER $\beta$  binds and regulates several important tumor suppressors and oncogenes in CRC cells, such as CST5 and LRP6, consistent with its proposed antitumorigenic activity. We also revealed the p65 cistrome in CRC cell lines



and showed that ER $\beta$  alters the p65 chromatin binding in a cell-type-dependent manner. We found that ER $\beta$  chromatin binding sites were enriched among circadian clock genes and also that ER $\beta$  modulates p65 binding to core clock genes in CRC cells, indicating potential cross-talk between ER $\beta$  and circadian clock gene regulation.

The final aim was to investigate the impact of ER $\beta$ , and estrogen signaling in high-fat diet (HFD) induced inflammation in colon, explored in paper III. We fed mice with an HFD for 13 weeks and treated them with estrogenic ligands for the last three weeks prior to sacrifice. The colon transcriptome showed predominant sex differences, and selective activation of ER $\beta$  reduced macrophage infiltration and epithelial cell proliferation induced by HFD. We demonstrated that ER $\beta$  opposes HFD-induced dysregulation of core circadian clock genes *in vivo*, further strengthening the role of ER $\beta$  in circadian rhythm.

Taken together, these results highlight the chemopreventive potential of ER $\beta$  in CRC in both sexes. The identified cross-talk with TNF $\alpha$ /NF $\kappa$ B pathway, Wnt signaling, regulating genes involved circadian clock, and tumorigenesis reflected ER $\beta$  protection/antitumor activity against CRC progression and development (as illustrated in Figure 1).




Figure 1. Schematic summary for the novel role of ER $\beta$  in the prevention of CRC.

## LIST OF SCIENTIFIC PAPERS

- I. Hases L, **Indukuri R**, Birgersson M, Nguyen-Vu T, Lozano R, Saxena A, Hartman, J, Frasor J, Gustafsson JÅ, Katajisto P, Archer A, Williams C. Intestinal estrogen receptor beta suppresses colon inflammation and tumorigenesis in both sexes. (2020) Cancer Letters. 492:54-62.
- II. **Indukuri R**, Jafferli MH, Song D, Damdimopoulos A, Hases L, Zhao C, Archer A, Williams C. Genome-wide estrogen receptor  $\beta$  chromatin binding in human colon cancer cells reveals its tumor suppressor activity. (2021) International Journal of Cancer. 1-15.
- III. Hases L, Archer A, **Indukuri R**, Birgersson M, Savva C, Korach-André M, Williams C. et al. High-fat diet and estrogen impacts the colon and its transcriptome in a sex-dependent manner. (2020) Scientific Reports. 10: 16160.
- IV. **Indukuri R**, Hases L, Archer A, Williams C. Estrogen Receptor Beta Influences the Inflammatory p65 Cistrome in Colon Cancer Cells. (2021) Frontiers in Endocrinology. 12:650625.



# CONTENTS

1	Introduction .....	1
1.1	Colorectal cancer.....	1
1.1.1	Epidemiology .....	1
1.1.2	Risk factors.....	1
1.1.3	Tumor staging and subtypes .....	2
1.1.4	Diagnosis and treatments .....	2
1.2	Pathways involved in CRC .....	3
1.2.1	Molecular pathogenesis.....	3
1.2.2	Wnt/ $\beta$ -catenin signaling pathway .....	4
1.2.3	RAS/MAPK pathway.....	5
1.2.4	p53 pathway .....	5
1.2.5	TGF- $\beta$ pathway .....	6
1.2.6	TNF $\alpha$ /NF $\kappa$ B signaling .....	6
1.3	Colorectal cancer - preventable cancer?.....	6
1.3.1	Inflammation .....	6
1.3.2	Inflammatory bowel disease .....	7
1.3.3	Obesity and diet .....	7
1.3.4	Sex and hormonal factors.....	8
1.4	Estrogens in CRC.....	8
1.4.1	Estrogens .....	8
1.4.2	Estrogen receptor alpha and beta.....	9
1.4.3	ER Structure .....	10
1.4.4	ER $\beta$ splice variants .....	11
1.4.5	GP130 .....	11
1.4.6	Ligands .....	12
1.4.7	ER signaling pathways .....	13
1.4.8	Enhancers and epigenetic mechanisms .....	15
1.5	ER $\beta$ role in CRC .....	16
1.5.1	Major challenges in the field.....	16
1.5.2	Epidemiological evidence .....	16
1.5.3	CRC suppression by ER $\beta$ .....	17
1.5.4	ER cross-talk .....	18
1.5.5	ER interactome studies .....	19
2	Aims.....	20
3	Material and Methods.....	21
3.1	Mouse models .....	21
3.1.1	ER $\beta$ KO Vil mouse.....	21
3.1.2	AOM/DSS-induced colitis .....	21
3.1.3	High-fat diet (HFD) induced colon inflammation.....	22
3.1.4	Tissue collection .....	22

3.2	Invitro studies .....	22
3.2.1	Colon cancer cell lines.....	22
3.2.2	Chromatin Immunoprecipitation .....	23
3.2.3	Luciferase assays .....	24
3.2.4	Statistical analysis.....	25
4	Results and discussion .....	26
4.1	Paper I: Intestinal estrogen receptor beta suppresses colon inflammation and tumorigenesis in both sexes.....	26
4.2	Paper II: Genome-wide estrogen Receptor $\beta$ chromatin binding in Human colon cancer cells reveals its tumor suppressor activity.....	28
4.3	Paper III: High-fat diet and estrogen impacts the colon and its transcriptome in a sex-dependent manner .....	30
4.4	Paper IV: Estrogen receptor beta influences the inflammatory p65 cistrome in colon cancer cells .....	32
5	Conclusions .....	35
6	Future perspectives .....	36
7	Acknowledgments .....	38
8	References .....	40

## LIST OF ABBREVIATIONS

5-ASA	5-aminosalicylic acid
5-FU	5-fluorouracil
AF1/2	Activation function 1/2
AOM	Azoxymethane
AP-1/2	Activator protein 1/2
APC	Adenomatous polyposis coli
AR	Androgen receptor
ARNTL	Aryl hydrocarbon receptor nuclear translocator like
ATAC-Seq	Assay for Transposase-Accessible Chromatin using sequencing
BCL3	B-cell lymphoma 3-encoded protein
BIRC3	Baculoviral IAP repeat-containing 3
BMI	Body mass index
BRAF	V-RAF murine sarcoma viral oncogene homolog B1
Co A	Co activator
CCND1	Cyclin D1
CD	Control diet
CEA	Carcinoembryonic antigen
ChIP	Chromatin immunoprecipitation
ChIP-Seq	Chromatin immunoprecipitation followed by sequencing
CIMP	CpG island methylator phenotype
CIN	Chromosomal instability
CK1	Casein kinase1
CMS	Consensus molecular subtypes
COX2	Cyclooxygenase2
CRC	Colorectal cancer
CREB	Cyclic AMP response element-binding protein
CRISPR	Clustered regularly interspaced short palindromic repeats
CTNNB	Beta-catenin
CTSD	Cathepsin D

DMEM	Dulbecco's modified eagle medium
DMSO	Dimethyl sulfoxide
DNA	DNA binding domain
DPN	Diarylpropionitrile
DSS	Dextran sodium sulfate
E1	Estrone
E2	17 $\beta$ -estradiol
E3	Estriol
E4	Estetrol
EGF	Epidermal growth factor
EGFR	Epidermal growth factor receptor
EMT	Epithelial-mesenchymal transition
ERE	Estrogen response element
ER $\alpha$ / $\beta$	Estrogen receptor $\alpha$ / $\beta$
ER $\beta$ KO <sup>vi</sup>	ER $\beta$ knockout mice
FAP	Familial adenomatous polyposis
FBS	Fetal bovine serum
FEB	Fecal occult blood
FOXA	Forkhead box a
FOXM1	Forkhead box protein M1
FOXO	Forkhead box O
GATA	GATA binding protein
GP <sup>ER</sup> 1	G protein-coupled estrogen receptor 1
GSK	glycogen synthase kinase3
H3K27 <sup>ac</sup>	Histone 3 lysine 27 Acetylation
H3K4 <sup>me</sup>	Histone 3 lysine 4 methylation
H&E	hematoxylin and eosin
HDI	Human development index
HFD	High-fat diet
HIC	Chromosome conformation capture (HiC)
HRT	Hormone replacement therapy
IBD	Inflammatory bowel disease

IgG	Immunoglobulin G
IHC	Immunohistochemistry
IL	Interleukin
JUN	Jun Proto-Oncogene
KLF	Kruppel-like factor
KRAS	Kirsten rat sarcoma viral oncogene
LBD	Ligand binding domain
LDL	Low-density lipoprotein
LEF	Lymphoid enhancer factor
LGR5	Leucine-rich repeat-containing G protein-coupled receptor 5
LRH1	Liver receptor homolog 1
LRP	Lipoprotein receptor-related protein
LRP5/6	Lipoprotein receptor-related protein 5/6
LXR	Liver X receptors
MAPK	Mitogen-activated protein kinase
MMR	Mismatch repair
MPA	Medroxyprogesterone acetate
mRNA	Messenger RNA
MSI	Microsatellite instability
MYB	Myeloblastosis viral oncogene
MYC	Master regulator of cell cycle entry and proliferative metabolism
NFκB	Nuclear factor-κB
NR	Nuclear receptor
NSAID	Non-steroidal anti-inflammatory drugs
NTD	N-terminal domain
PCR	Polymerase chain reaction
PER2	Period circadian protein homolog 2
PPAR	Peroxisome-proliferator-activated receptors
PPT	Propylpyrazoletriol
PROX1	Prospero homeobox 1
RAR	Retinoic acid receptors



RIME	Rapid immunoprecipitation mass spectrometry of endogenous proteins
ROS	Reactive oxygen species
RUNX	Runt-related transcription factor 1
RXR	Retinoid x receptor
SERM	Selective estrogen receptor modulator
SFREs	SF-1 response elements
SMAD	Mothers against decapentaplegic homolog 2
SP-1	Stimulating protein 1
SRC-1	Steroidal co-receptor activator 1
TCF	T cell factor
TF	Transcription factor
TGF $\beta$	Transforming growth factor beta
TGF $\beta$ R1/2	Transforming growth factor beta receptor 1/2
TNBS	Trinitrobenzene sulfonic acid
TNF $\alpha$	Tumor necrosis factor $\alpha$
TNM	Tumor-node-metastasis
TP53	Tumor protein p53
TRE	TPA response element
VDR	Vitamin d receptor
WNT	Wingless-type MMTV integration site family
WT	Wild type

# 1 INTRODUCTION

Globally, we have entered an era of increased living standards and affordable access to quality healthcare with better diagnosis methods and treatment of diseases. Life expectancy is also raised in most parts of the world as a result of access to health care. Improved medical treatments, however, have decreased death rates primarily from communicable diseases. By contrast, cancer-related deaths have increased by 40% over the past 40 years. A further 60% increase in cancer deaths was expected by 2030, with an estimated 13 million people dying of cancer, as reviewed in (4, 5). However, a recent study in the US population reported a 26% decline in the cancer death rates over two decades (6). Colorectal cancer (CRC) has become a leading cause of cancer, affecting 1.23 million patients each year and accounting for 10% of cancer-related deaths in Western countries (4, 5).

## 1.1 Colorectal cancer

CRC is defined as a tumor that originates from the colon and rectum. It is the third most prevalent cancer in the world, with an estimated 880 792 deaths each year, 1.8 million new cases, and 1.2 million deaths expected in the next ten years (7, 8). Roughly 66% of the cases are colon cancers, and the remaining are rectal cancers (9). After lung, breast, and prostate cancer, CRC is the fourth most common malignant tumor in terms of incidence, yet it is ranked the third in regards to cancer mortality (8).

### 1.1.1 Epidemiology

Colorectal cancer classifies as a symbol of socioeconomic growth, and, in nations experiencing significant development, prevalence rates are inclined to increase consistently with a high human development index (HDI) (10, 11). An upsurge in previously low-risk and lower HDI nations signifies lifestyle changes along with diets, such as changes toward an amplified consumption of animal-source diets, low physical activity, and body mass is independently correlated with CRC risk (12, 13).

For decades, CRC was considered an old age disease; however, recent data indicate increased CRC incidence in young adults under the age of 50 (14-16). Most of these individuals under the age of 50 would not meet the screening criteria. One recent study identified the rise of rectal cancers compared to colon predominantly in young individuals (14). The rising incidence in young birth cohorts indicates the influence of dietary patterns, excess body weight, and lifestyle factors (13).

### 1.1.2 Risk factors

CRC is a complex disease being influenced by multiple factors such as aging, chronic inflammation, poor dietary habits, obesity, and lifestyle. The most significant risk factor for CRC is older age; the disease is rare before age 40 (17). Around 80% of CRC cases are sporadic; only 10% are caused by genetic predisposition such as Familial adenomatous polyposis (FAP) and Lynch Syndrome. FAP is caused by inherited germline mutations in the

tumor suppressor gene adenomatous polyposis coli (APC). Lynch syndrome is characterized by mutations in mismatch repair genes (MMR) (18, 19).

Male gender increases the risk of CRC incidence and overall survival (17). Patients with inflammatory bowel disease and ulcerative colitis also have a higher risk of developing colon cancer (20). In CRC, substitution mutations that change single amino acids were the highest number of mutations indicating CRC's are most vulnerable to environmental factors (21). Thus, lifestyle-related diseases, such as metabolic syndrome, cause between 40-70% in terms of CRC development, and diet has been shown to influence CRC risk (22-24). A recent study indicated that high intakes of red meat and processed meat increase the risk of CRC (25), while phytoestrogens or isoflavones dietary intake provided a protective effect in CRC patients (26). High vitamin D levels are correlated with improved survival and decreased recurrence. Supplementary calcium intake was associated with reduced colon cancer risk (27, 28). In the UK population, using a modeling approach, 5% of diagnosed colon cancer cases were estimated due to low levels of physical activity. A meta-analysis showed that high physical activity reduces the overall risk of CRC by 24% compared to the least active individuals (29, 30).

### **1.1.3 Tumor staging and subtypes**

In colorectal cancer, the tumor's pathological stage is generally considered the most critical determinant of outcome. The TNM classification is the most widely used, which considers tumor invasion, spread to lymph nodes, and metastasis to stage CRCs (31). The high degree of heterogeneity in CRC, at both genomic and transcriptomic levels, leads to further categories based on gene expression profiles.

Consensus Molecular Subtypes (CMS) classification identified four subtypes (CMS1-CMS4) and helped in personalizing oncological care by distinguishing the patients who may benefit from adjuvant chemotherapy (32). The CMS1 subtype is associated with major histological changes, more often seen in women, and is linked to MMR status. The tumors of the CMS2 are the most diverse among all, whereas the CMS3 subgroup contains a high incidence of KRAS mutations. The CMS4 group is considered a pro-inflammatory type due to increased mesenchymal and stromal genes instead of genes involved in cancer, indicating the importance of inflammation in CRC. The CMS4 group had the most advanced stages of CRC of all the groups, as well as the highest tumor budding compared to the CSM2, CSM3 subtypes(33).

### **1.1.4 Diagnosis and treatments**

CRC's routine diagnosis is based on physical examinations, colonoscopy, and computer tomography colonography. The serum tumor marker carcinoembryonic antigen (CEA) or fecal occult blood is also measured using laboratory tests (FEB). In most of Europe, Canada, and North America, screening initiatives testing FEB and using colonoscopy are currently in progress to detect CRC early (34). A CRC screening program started in Stockholm in 2008 and

within five years identified 0.1 percent with CRC out of 1.8 percent tested with a colonoscopy (35).

Studies have been conducted using non-steroidal anti-inflammatory drugs (NSAIDs) to prevent CRC. NSAIDs such as aspirin and celecoxib inhibit the cyclooxygenase-2 (COX-2) enzyme, which induces inflammation and reduces the risk of colon cancer. Two extensive clinical studies revealed that long-term use of NSAIDs reduces CRC incidence by 37% (36, 37). However, these clinical trials stopped early due to cardiac side effects of COX-2 inhibitors such as celecoxib and rofecoxib (17, 38, 39). A randomized control trial of 164,225 patients confirmed that aspirin increased the risk of bleeding (40). Hence COX-2 inhibitors, which had previously been approved by the European Union (41) and the United States (42), were no longer used to prevent CRC, and there is a need for developing new and safe targets.


The cure for colorectal cancer is determined by the degree to which cancer has spread. Complete surgical resection is the only approach in treating colorectal cancer in stages between I and III. Primarily, 2 out of 3 individuals who go through surgical resection will have cancer regrowth or metastases. Thus, to treat and prevent a recurrence, chemotherapy treatment with single drug 5-Fluorouracil (5-FU) or in combination with oxaliplatin, capecitabine used (43).

## **1.2 Pathways involved in CRC**

Notably, the three common pathways in CRC development are chromosomal instability (CIN), microsatellite instability (MSI), and CpG island methylator phenotype (CIMP). Primarily, the CIN pathway accounts for 85% of the sporadic CRCs. It is mainly caused by chromosomal imbalance and loss of function mutations in tumor suppressor genes. Fundamentally, MSI is categorized by mutations in MMR genes regulating microsatellites in DNA replication (44). MSI creates frameshift mutations that inactivate genes involved in the MMR process (45). Aberrant methylation of CpG islands leads to epigenetic instability in CRC, and 10-20% CRCs contain a high percentage of methylated CpG loci (46). Nonetheless, the mechanisms leading to CIMP are still unknown.

### **1.2.1 Molecular pathogenesis**

The evolution of colon cancer is mainly due to the progressive accumulation of genetic mutations through which cells acquire the ability to sustain cancer hallmarks. The six biological hallmarks of cancer are proliferative signaling, inhibiting growth suppressors, resisting cell death, evading apoptosis, and inducing angiogenesis. Genetic and epigenetic alterations activate oncogenes and inactivate tumor suppressor genes to enable cancer hallmarks (47). Aberrant colonic crypts drive the normal colonic epithelium into adenomatous polyps, benign tumors with malignant potency. Benign adenomas develop into carcinomas over 10-15 years due to the accumulation of genetic mutations primarily in adenomatous polyposis coli (APC), KRAS proto-oncogene, and tumor protein p53 (TP53) genes. The pathways involved in the molecular progression of CRC are depicted in figure 2 and discussed as follows.



**Figure 2:** Molecular progression of colorectal cancer. Inspired by Kuipers et al., 2015.


### 1.2.2 Wnt/ $\beta$ -catenin signaling pathway

Wnt signaling is a central mechanism that controls cell fate decisions which mainly acts through frizzled protein, low-density lipoprotein receptor-related protein 6 (LRP6) receptors. In a normal colon, the Wnt ligand maintains stem cell homeostasis, and the Wnt signaling target gene *Lgr5* is a well-established marker for intestinal stem cells (48). CRC cells with high expression of *Lgr5* clonally expand and form colonies compared to cells without *Lgr5* expression (49). High Wnt activity was observed in colon cancer stem cells and cancer stemness regulation (50).

Wnt signaling is divided into two pathways, canonical, dependent on the  $\beta$ -catenin and, non-canonical pathway, independent of  $\beta$ -catenin.  $\beta$ -catenin is encoded by the gene *CTNNB1* and plays a vital role in cell morphology. Both cytoplasmic and nuclear  $\beta$ -catenin levels are strictly regulated by a group of proteins called  $\beta$ -catenin destruction complex, which keeps the levels of  $\beta$ -catenin low in the absence of a Wnt signal. This complex is formed by scaffolding protein axin, glycogen synthase kinase 3 (GSK3), casein kinase 1 (CK1), and APC protein. APC is a tumor suppressor gene that negatively regulates  $\beta$ -catenin and controls proliferation, differentiation, and migration promoted by Wnt signaling. APC is mutated in more than 80% of CRC patients (51). When Wnt binds to the frizzled receptors and LRP5/LRP6 in the canonical pathway, it inactivates the  $\beta$ -catenin destruction complex and activates  $\beta$ -catenin.  $\beta$ -catenin, which does not possess a DNA-binding domain, interacts with TCF (T-cell factor) or LEF (lymphoid enhancer factor) transcription factors to regulate Wnt target genes involved in cell migration, proliferation, and survival such as *CCND1* (cyclin D1), *PTGS2* (COX-2) and *MYC* (c-Myc) (52).

In the absence of a Wnt signal,  $\beta$ -catenin destruction complex phosphorylates  $\beta$ -catenin, while APC constantly ubiquitinates and degrades the  $\beta$ -catenin. The central region of the APC protein mainly coordinates with  $\beta$ -catenin through seven 20 amino acid repeats. APC mutations in colon cancer were primarily found in the third 20 amino acid repeat of the protein, which has been shown to be the most highly phosphorylated site with an increased affinity for  $\beta$ -catenin (53). APC gene inactivation fails to degrade  $\beta$ -catenin.  $\beta$ -catenin then increases transcription through TCF (Figure 3). Inactivating APC gene mutations initiate the majority of CRCs by

increasing Wnt signaling. Mutations of the APC gene have been reported as an early CRC mutation for the acquisition of cancer hallmark inactivated tumor suppressor genes (54). In FAP patient's germline APC mutations causes hundreds of polyps in the colon and rectum (18).



**Figure 3.** Wnt signaling pathway in colorectal cancer (Pino et al. Gastroenterology 2010, permission obtained to print 5055420182095)

### 1.2.3 RAS/MAPK pathway

Adenoma to carcinoma progression is initially influenced by the KRAS gene, a proto-oncogene encoding for an ATPase protein that transmits an extracellular signal through MAPK and enhances cell proliferation. Activating mutations in the KRAS gene permanently activates the protein to sustain proliferative signaling and avoid apoptosis. Single point mutations in codon 12 and 13 lead to an oncogenic phenotype and are hotspot mutations in the KRAS gene (55).

### 1.2.4 p53 pathway

p53 is a tumor suppressor that regulates cell cycle arrest and apoptosis. Inactivating p53 mutation is a common and critical molecular step in CRC progression. Loss-of-function p53 mutations are most common (56, 57) and result in proliferative activity and resistance to cell death. Unlike mutations of the APC and KRAS genes, and unlike some other cancers (such as non-melanoma skin cancers), p53 mutations usually occur in the later stages of colorectal carcinogenesis.

### **1.2.5 TGF- $\beta$ pathway**

TGF- $\beta$  is a member of the cytokine family regulating epithelial systems and inflammatory processes, particularly in the gut. TGF- $\beta$  receptors (TGFR1, TGFR2) phosphorylate SMAD2 and SMAD3, which then form a complex with SMAD4 and move to the nucleus to regulate transcriptional activity. TGF- $\beta$  also maintains tissue homeostasis and regulates cell proliferation, adhesion, and survival. Cancer cells escape this pathway during malignant evolution through mutations that truncate TGF- $\beta$  receptors and inactivate the SMAD4 gene (58, 59).

### **1.2.6 TNF $\alpha$ /NF $\kappa$ B signaling**

The function of the nuclear factor of  $\kappa$ B (NF $\kappa$ B) was previously thought to be immune-related, but oncogenic mutations activated NF $\kappa$ B in lymphoid cancers have revealed its important role in cancer. NF $\kappa$ B is a family of transcription factors, which regulate innate immune and inflammatory responses. A constitutive expression of NF $\kappa$ B in cancers revealed other functions of NF $\kappa$ B, such as cell proliferation, migration, and apoptosis (60, 61). There is a large amount of evidence suggesting NF $\kappa$ B participation in carcinogenesis by activating cellular gene expression. To date, five different NF $\kappa$ B family members have been identified; NF $\kappa$ B1 (p50/p105), NF $\kappa$ B2 (p52/p100), RelA (p65), RelB, and c-Rel. NF $\kappa$ B protein dimers bind to the promoter of target genes on a standard specific sequence known as " $\kappa$ B site." The term NF $\kappa$ B mostly refers to p65-p50 heterodimer, which is the primary dimer in most cell types. NF $\kappa$ B transcriptional activity is dependent mainly on p65, which contains the most potent transcriptional active domain (61-64).

## **1.3 Colorectal cancer - preventable cancer?**

As described above, the origins and causes of colorectal cancer (CRC) are many. Many risk factors are modifiable (17), which allows the potential prevention of most CRCs development (22-24).

### **1.3.1 Inflammation**

Inflammation is one of the most important hallmarks of cancer, promoting cancer development and progression through different mechanisms (65). Inflammatory diseases such as colitis, pancreatitis, and hepatitis, have been associated with an increased risk of colon, pancreatic, and liver cancers, respectively. A prolonged inflammatory condition attracts macrophages, leukocytes which secrete tumor necrosis factor- $\alpha$  (TNF $\alpha$ ), and reactive oxygen species (ROS). This aggravates DNA damage (66) and results in mutations (67, 68). Further, chronic inflammation profoundly affects metastasis through epithelial-mesenchymal transition (EMT), providing mobility and the ability for cells to migrate from the primary tissue (69). Through activation of wound healing processes, the inflammatory response augments angiogenesis and invasion, advancing the tumors' development (70).

As an inhibitor of cyclooxygenase-2 (COX-2), a component that ranks as an important mediator of the inflammatory reaction, Aspirin was shown to be substantially effective in preventing adenomatous polyp and CRC (71, 72). However, Aspirin is not advisable because of the increased risk of bleeding. Frequent usage of Non-Steroidal Anti-Inflammatory Drugs (NSAID) is associated with a lower risk of CRC (71, 72), even though the case is different when there is an occurrence of BRAF mutation-caused CRC (73). However, trials with NSAID celecoxib have ceased due to cardiac side effects (38, 40). New treatments are necessary due to the risks posed by current prevention therapies.

### **1.3.2 Inflammatory bowel disease**

Inflammatory bowel disease (IBD) results from the chronic inflammation of the intestine triggered by environmental and immunoregulatory factors, unbalanced gut microbiota, and genetic predisposition. IBD has increased rapidly worldwide and also, 25% of the detected cases were below the age of 20, indicating a rise in children and adolescents (74). IBD is categorized into Crohn's disease and ulcerative colitis. In many cases, Crohn's disease has its effects on the entire GI tract, whereas ulcerative colitis can be termed as a disease of the large intestines, specifically the sigmoid and rectal parts (75). IBD is a chronic condition that can be clinically managed using 5-aminosalicylic acid (5-ASA), doses of immunomodulatory (azathioprine), and anti-TNF $\alpha$  drugs such as infliximab. Nonetheless, surgery may be preferred in certain cases (74).

IBD increases the risk of CRC, and eventually, 20-30% of IBD patients will develop colitis-related colorectal cancer (CA-CRC) (76, 77). CA-CRC shows a complex malignancy by acquiring p53 mutations at an early stage compared to the sporadic type of CRC, where APC mutations are the first events (78). Sex differences were identified in CRC development in patients with IBD; men with IBD are at 60% increased risk to develop CRC compared to women (20).

### **1.3.3 Obesity and diet**

Epidemiological evidence suggests that having a high Body Mass Index BMI (BMI) raises the risk of developing CRC (79). A large cohort study conducted among ~80,000 females identified that being overweight was connected to an increased risk of early-onset CRC in women. The foremost aspects of obesity are chronic low-grade inflammation and excess adipose tissue, which enhances the release of free fatty acids, interrupting insulin signaling in the skeletal muscle (80). Also, HFD is illustrated to affect intestinal permeability, microbiota, along with inflammation (81-83). A new study suggests that high-fat diet advances CRC development upsets the balance of bile acids within the intestine, activates a hormonal signal that helps cancerous cells thrive. In this case, such findings could clarify why colorectal cancer, a disease that can take years to develop, is becoming more common in younger people, following current trends indicating that higher-fat diets are more prevalent. (84). Consumption of an HFD causes metabolic dysfunction by disrupting the systemically harmonized circadian rhythm (85). For instance, a high-fat diet modifies the food intake of rats with extra food intake



during the day (86). It disrupts the rhythmicity of *Clock*, *Bmal1*, and *Per2* within the liver as well as the adipose tissue (87) (88). In the gut, circadian rhythm plays an essential role in regulating intestinal permeability, cell proliferation, gut microbiota, and metabolism, and perturbation of this fine balance can result in IBD and CRC (as reviewed in (89)). The dysregulation of circadian genes results in increased pro-inflammatory cytokines plus the body adiposity regarding the diet-induced obese mice (86). Indeed, meta-analyses have discovered a relationship between night shift work, obesity, and augmented CRC risk (90, 91).

### **1.3.4 Sex and hormonal factors**

There are apparent sex differences in CRC. Compared to women, the prevalence of CRC is significantly higher in men. More aggressive proximal CRC is most common in women, while men predominantly suffer from distal CRC (as reviewed in (92)). Studies have been conducted to showcase the varying origins and causes of both distal and proximal CRC. Men and women have different gastrointestinal transit times, impacting the interaction between anti-carcinogenic, carcinogenic agents and colonic mucosa. Abnormal glucose metabolism promotes the growth and progression of CRC as reviewed in (93, 94), and sex differences highly influence glucose metabolism and DNA methylation patterns which alter insulin production (95). Sex hormones also play an essential role in the progression of certain cancers. Estrogens also impact gastrointestinal tract diseases (96, 97), including gastroesophageal reflux disease (98), irritable bowel syndrome, and IBD (99).

## **1.4 Estrogens in CRC**

### **1.4.1 Estrogens**

Estrogens are steroid hormones that are regarded as necessary for sexual and reproductive development in women and men. Estrogens are involved in various physiological functions within the female body, such as bone integrity, muscle mass, subcutaneous visceral fat, circadian rhythm, and homeostasis (100, 101). In men, estrogens are also critical during the development of their reproductive organs and sex differentiation of the brain (102).

The term estrogen denotes several chemically similar hormones within this group which include estrone (E1), 17 $\beta$ -estradiol (E2), estriol (E3), and estetrol (E4) (103). E2 is the most predominant circulating form of estrogen in pre-menopausal women and is 10 times more potent than E1 and 80 times more potent than E3 (104). The primary production of E2 is in the ovaries of premenopausal women. It is synthesized in the granulosa cells of the ovaries through a process called steroidogenesis. Dietary cholesterol, particularly low-density lipoprotein (LDL)-cholesterol, is the primary substrate for steroid hormone biosynthesis (103). Estrogens are a result of the action of aromatase, responsible for the conversion of testosterone to estradiol and androstenedione to estrone. The granulosa cells of the ovarian follicles and the corpora lutea produce E2. The synthesis of estrogens also occurs in smaller amounts in other tissues that express aromatase, such as the adrenal glands, liver, pancreas, adipose tissue, and breast (106).

In males, E2 is produced through aromatization in Sertoli cells, Leydig cells, and mature spermatocytes of the reproductive tract (102). In men and postmenopausal women, E1 is the most prominent estrogen and comes from a local synthesis of estrogens in extra gonadal target tissues by aromatization (105). Through this, the E1 may also be converted into E2 in peripheral tissues. These include adipose and breast tissue, vascular endothelium, smooth muscle cells, brain tissue, and bone cells (105, 107). The estrogens local production implies the signaling modality from endocrine to act locally as a paracrine or intracrine factor (105). E3 and E4 are only synthesized during pregnancy by the placenta and fetal liver, respectively (108, 109). All estrogens can bind to nuclear and membrane estrogen receptors, but their affinity and strength of the response vary. The most potent among all estrogens in inducing cell proliferation is E2, followed by E1 and E3 (110).

A recent study conducted using gastric biopsies found that parietal cells produce large amounts of estrogens within gastric mucosa and eventually secrete them to the portal vein (111). It is hypothesized that gastric E2 binds to the ER and controls certain functions or is metabolized into inactive estrogens by liver (112). Thus, it explains the pathophysiological importance of estrogens in gastrointestinal tract.

#### **1.4.2 Estrogen receptors alpha and beta**

Estrogen receptors (ERs) are the mediators of estrogens (102, 103). Estrogen mainly acts through three receptors, estrogen receptor alpha (NR3A1/ESR1), estrogen receptor beta (NR3A2/ESR2), and G protein-coupled estrogen receptor (GPER1, see below). ER $\alpha$  and ER $\beta$  are nuclear receptors. Nuclear receptors (NRs) are multifunctional proteins transducing signals of their cognate ligands. Humans have 48 nuclear receptors, including steroid, thyroid, vitamin D, and retinoid receptors. These are involved in signal transduction and act as ligand-inducing transcriptional regulators which control the activity of specific gene networks. They can be divided into three categories: endocrine NRs, orphan NRs and adopted NRs based on ligand binding. Endocrine NRs rely on hormones as their endogenous ligands, such as ERs. Orphan NRs have no known natural ligand, while adopted NRs have their natural ligand discovered after previously being considered orphans (113). The various forms of NRs include monomers (steroidogenic factor-1 (SF-1), estrogen-related receptor  $\beta$ ) (114) and homodimers (ERs, and androgen receptors (AR), heterodimers with other nuclear receptors (retinoid X receptor (RXR), thyroid receptor, and vitamin D receptor (VDR), among others (115).

The most researched NR functions are the ones of AR in the prostate and ER in the breast. AR is a steroid receptor for testosterone which is a critical driver in prostate cancer, and several anti-androgen drugs are in use to reduce AR activity (116). ER $\alpha$  initiated and proliferate breast cancer, and a majority of the breast cancers contain ER $\alpha$  expression and responds to hormone treatment (117). About 70% of NR binding sites were prominent in open chromatin (113, 118). Nevertheless, NRs recruit a variety of chromatin-modifying factors to facilitate binding to target sites within closed chromatin.

HNF4, ER, LRH1, PPAR, VDR, RXR, RAR and LXR are the nuclear receptors expressed in the colon (113, 119). In both mouse and human models of CRC, the enteric NR transcriptome is downregulated during tissues' progression from normal intestinal epithelia to dysplastic lesions. Therefore, a therapeutic, diagnostic potential is proposed for these transcription factors in CRC (120, 121).

ER $\alpha$  was discovered in 1968 (122), and after 27 years, the second distinct estrogen receptor ER $\beta$  was identified in rat prostate by the Gustafsson group. The ER $\alpha$  and ER $\beta$  genes are located on different chromosomes (6q25.1 and 14q22-24) (123-125). ER $\alpha$  is mainly expressed in the endometrium, ovary, mammary gland, bone, uterus, smooth muscles, and epididymis within the human body. On the other hand, ER $\beta$  can be found in the ovaries, testis, adrenal gland, and spleen (126).

### 1.4.3 ER structure

All nuclear receptor proteins are composed of a variable N-terminal domain (NTD), a DNA binding domain (DBD), a hinge region, a conserved ligand-binding domain (LBD), and a variable C-terminal domain. The DNA binding domain has two zinc finger motifs that serve as hooks and facilitate binding to chromatin in the nucleus. Each class has its own DNA binding recognition sequences, which is a distinguishing feature. Nuclear receptor ligand-binding domains are functionally quite similar, but their specificity and affinity for particular ligands differ (127).

Estrogen receptors are composed of five functional domains. Beginning from NH<sub>2</sub>- to COO-terminus, Activation function 1 (AF1) and NH<sub>2</sub> terminal domain (NTD) forms domain A/B, DNA binding domain (DBD) forms domain C, domain D is hinge region, ligand binding domain (LBD), and Activation function 2 (AF2) located within domain E and domain F in C-terminal of protein.

The NTD/AF1 domain is capable of ligand independent activity through phosphorylation and subsequent transcriptional activation of target genes (128, 129). DBD recognizes specific DNA sequences denoted as estrogen response elements (ERE) in target genes, facilitating ER binding to stimulate or repress the transcription of these genes (130). LBD/AF2 domain contains a hormone-binding pocket that facilitates ligand binding and dimerization. The orientation of the AF2 domain determines the functions activated by ligands as an agonist or antagonist (131). Besides, AF1 and AF2 are important regions that activate transcription and interact with other nuclear receptors or co-activators (132). The DBD and LBD are connected by the D domain, which includes amino acids that are important for post-translational modification (133). Additionally, the F domain controls the transcription of genes in a cell specific and ligand independent way (134, 135).


DNA-binding domains of ER $\alpha$  and ER $\beta$  are highly conserved, but the two nuclear receptors differ significantly in their N-terminal domains and in their ligand-binding domains. At hinge regions of ER $\alpha$  and ER $\beta$ , they share only 36% homology (124). Although both receptors

strongly bind estrogen, the overall homology of their ligand-binding domains is less than 55%, which has enabled the development of receptor-selective ligands (124, 134, 136).

#### 1.4.4 ER $\beta$ splice variants

*ESR2* contains eight encoding exons, which spans 254 kb, and it has several isoforms due to alternative splicing of the last coding exon (137). These isoforms are modified in the 3' end of hER $\beta$ 1 (wild type ER $\beta$ ) (Figure 4). Of all, wild-type ER $\beta$  (530 aa) is the only isoform that binds to the ligand (7). ER $\beta$ cx can heterodimerize with ER $\alpha$ , hence widely seen as a dominant-negative receptor for ER $\alpha$  (138, 139). It has since been established in a recent study that ER $\beta$  splice variants ER $\beta$ cx and ER $\beta$ 5 are present in granulosa cell tumors (140). The study, however, does not clearly outline their roles in the pathogenesis and progression of cancer (137, 140).

There is evolutionary conservation of ERs between species (141). Rodents have two known ER $\beta$  splice variants: a full-length rat and mouse ER $\beta$  WT isoforms (549 amino acids), with a sequence similarity of 99% (131) and both mouse and rat ER $\beta$ 2 isoforms with an 18-amino acid insertion in the LBD that leads to decreased ligand binding affinity of estrogen (142-144).



**Figure 4.** Estrogen receptor beta contains five domains, and all splice variants of ER $\beta$  identified in humans vary in F domain and the structural similarities between ER $\alpha$  and ER $\beta$ .

#### 1.4.5 GPER1

The third estrogen receptor is GPER1 (G protein-coupled receptor 1) (145). This receptor does not share structural similarities with ER $\alpha$  or ER $\beta$  and is not a transcription factor (146, 147). The G-protein coupled receptors (GPCRs) family entails the biggest cell surface receptor and contains seven transmembrane domains (148). Compared to other estrogen receptors, GPER1 has a poor binding affinity to ligand E2 (17 $\beta$ -estradiol) and requires high levels of ligand to become activated (149, 150). However, once activated by E2, GPER1 produces rapid responses and activates intracellular signaling cascades by increasing cAMP production (151) and intracellular calcium release (149). Specifically, estrogen activated GPER1 is involved in

various physiological functions in human tissues and organs, such as nervous, immune, reproductive, cardiovascular mechanisms, and cancer development and metastasis (152). The rest of this thesis will not focus on GPER1, and ER will refer to ER $\alpha$  and ER $\beta$ .

#### 1.4.6 Ligands

Estrogen is an endogenous ligand that activates ERs. Nevertheless, numerous other compounds presenting estrogenic effects that structurally or functionally resembles mammalian estrogens are recognized. Such compounds include phytoestrogens (naturally synthesized by plants) (153) and xenoestrogens (synthetic chemical compounds) from medicinal drugs, food additives, and plastics or environmental contaminants (154-156). Phytoestrogens can activate both ER's. Some phytoestrogens such as genistein, coumestrol, and liquiritigenin are more selective to ER $\beta$  based on their capability to bind to LBD with a higher affinity in comparison to that of ER $\alpha$  (157-159).

Whether natural or synthetic, ligands may be categorized as selective estrogen receptor modulators (SERMs), agonists, or antagonists. Agonistic ligands bind to ERs and trigger them, whereas antagonists block ERs activity upon binding. Agonists and antagonists bind at the exact position of the ERs LBD with different binding alignments. Agonist rearranges the ligand-binding domain conformation, creating helix 12 to rotate in a way to enable the free recruitment of co-activators while at the same time removing the co-repressors. By contrast, the antagonist disturbs helix 12 conformations to hinder association with co-activators (160, 161).

SERMs are a collection of synthetic nonsteroidal compounds that act as both agonists and antagonists of ERs, depending on the tissue type or estrogen receptor type. SERMs display antagonism in breast and uterus and a more agonistic nature in bones and liver (162). For this reason, they are utilized for the management of ER $\alpha$ -positive breast cancer (antagonism) and the deterrence of osteoporosis after menopause (agonism). A classic example is tamoxifen which acts as an antagonist in breast but as an agonist in uterus (163). Consequently, even though tamoxifen is the favored drug for treating ER-positive breast cancer, it can also prompt endometrial cell development and increase endometrial cancer risk (164, 165).

Another group of SERMs binds to only one of the ER subtypes. The ER $\alpha$ -selective agonist 4,4',4''-(4-propyl-1H-pyrazole-1,3,5-triyl) trisphenol (PPT) and the ER $\beta$ -selective agonist diarylpropionitrile (DPN) were used in the studies presented in this thesis. Notably, PPT binds to ER $\alpha$  with a 410-fold selectivity over ER $\beta$  (166), whereas DPN binds to ER $\beta$  with 70-fold selectivity over ER $\alpha$  (167).

To circumvent the classic actions of estrogens, such as stimulation of uterine proliferation, pathway-selective estrogen receptor (ER) ligands such as WAY-169916 were developed (168). WAY-169916 has shown strong anti-inflammatory activity in synoviocytes obtained from arthritis patients (169). Numerous other non-estrogenic compounds like androstane-3 $\beta$  and

17 $\beta$ -diol (3 $\beta$ D) have been shown to interact with the ERs (170, 171). One recent study also identified ketamine and its metabolites as a regulator of ER $\alpha$  (172).

#### **1.4.7 ER Signaling pathways**

ERs trigger transcriptional procedures and signaling events through direct binding to estrogen response element (ERE) in gene promoters or enhancers (genomic effects) or through processes that do not comprise a direct binding to the DNA (non-genomic effects), Figure 5 (103).

##### **Direct genomic signaling: ERE-dependent nuclear signaling**

In the classical pathway, ligand-activated ERs translocate into the nucleus, dimerizes, and binds directly to the consensus ERE sequence or estrogen half-sites. ERE is a 13-bp palindrome with a 3-bp spacer (5'-GGTCAnnnTGACC-3'), and ERE half-sites are either the proximal or distal half of the palindrome (173, 174). However, certain estrogen-responsive genes comprise EREs that diverge from the consensus through one or more nucleotides as well as being less potent regulators of the transcription than the consensus ERE (175). Nevertheless, regarding half ERE sites 5'-TGGGCTCA-3', controversy arises about whether ER can bind to these sites (176). A study that incorporated genetic deletions within the ER-bound enhancers by utilizing CRISPR in MCF7 cells showed that two enhancers containing half EREs could not compensate for the full ERE site and indicate that the position of full ERE immensely affects the enhancer activity (177).

It can be noted that homodimers (ER $\alpha$ -ER $\alpha$  or ER $\beta$ -ER $\beta$ ) and heterodimers (ER $\alpha$ -ER $\beta$ ) can be formed by ERs based on the tissue type and ligand. Even though ER $\alpha$  and ER $\beta$  selectively bind to EREs with similar affinity *in vitro*, the mechanism and degree of transcription that the ERs control significantly differs via the ERE-dependent signaling pathway (173, 178). *In vitro* gel shift assays have shown that compared to ER $\alpha$ , ER $\beta$  has lower binding to non-consensus EREs, which account for most estrogen-responsive elements (e.g., FOS, JUN, pS2, CTSD) (179).


##### **Indirect genomic signaling: ERE-independent nuclear signaling**

Unlike the classical pathway, countless ER binding sites lack ERE components (180-182); nonetheless, ER's control the transcription of these genes through an approach called transcription factor cross-talk (183-186). As stated by the most current studies, a projected 35% of genes that are targeted by estrogen do not contain EREs (103). Previous studies (187-190) have revealed the potential of ER to coact or tether with a range of coactivators, for instance, activating protein-1 (AP-1), stimulating protein 1 (Sp1), RUNX1, and NF $\kappa$ B. A cross-talk between ER $\alpha$  and NF $\kappa$ B has been studied. A few studies (191, 192) report that ER $\alpha$  represses NF $\kappa$ B activity, but on the contrary, other studies report that ER $\alpha$  enhances NF $\kappa$ B activity (193, 194) using the same cell lines.

The most widely studied ER coactivator is AP-1. ER bind members of AP-1 complex (Jun/Fos heterodimer) transcription factors. Indeed, mapping ER binding sites across the genome shows much overlap with AP-1 binding sites (195). In cell lines, when treated with E2, ER $\alpha$  and ER $\beta$

showed opposite transactivation of the AP-1 site: E2 activated transcription via ER $\alpha$ , while inhibited via ER $\beta$  (196). This was also noted in cyclin D1 gene expression, where E2-bound ER $\beta$  reduced expression of cyclin D1 and further blocked ER $\alpha$ -mediated activation of cyclin D1, as reviewed in (103).

Additionally, several estrogen-responsive genes lacking full EREs have ERE half-sites or binding elements for the orphan nuclear hormone receptor SF-1 [SF-1 response elements (SFREs)], serving as direct ER binding mechanisms (197). Several studies indicate that ER $\alpha$ , but not ER $\beta$ , binds to SFREs (198) and transcriptionally stimulates the SP1 component (199). Notably, studies point out many dissimilarities in ER $\beta$  and ER $\alpha$  coordinating with cofactors.



**Figure 5.** Different modes of ER signaling, including genomic, nongenomic, and signaling through growth factors. Inspired by Morselli et al., 2018.

### Indirect signaling: membrane receptors, mitochondrial events, signaling molecules

The ERs exert cell-specific immediate effects by activating signaling molecules such as the Ras/MAPK pathway and cAMP. Activated kinases phosphorylate the ERs in the N-terminal domain, causing receptor dimerization, binding to DNA, and initiation of gene expression in a ligand independent fashion (131, 200). Furthermore, estrogen that binds to GPER1 also stimulates estrogen-dependent stimulation of adenylyl cyclase and epidermal growth factor receptor (EGFR). Also, methylation, acetylation, ubiquitination, and sumoylation can restrain ER action via multiple mechanisms (201).

The rapid effects of estrogen, such as increased cAMP concentration in minutes (without gene activation), suggest that ERs confined within the plasma membrane are accountable for this action (202, 203). Some studies indicated that post-translational modifications such as palmitoylation increase the receptor's association with the plasma membrane. In the plasma membrane, E2 drives cell survival through ER $\alpha$  non-genomic signaling and cell death through ER $\beta$  non-genomic signaling (204). Additional to the plasma membrane, ERs have been reported in organelles such as mitochondria (205) and the endoplasmic reticulum (149, 206). However, ERs non-genomic activity is not as well understood as the genomic pathways, and additional research is required in this area.

#### **1.4.8 Enhancers and epigenetic mechanisms**

ERs primarily control gene expression by binding directly to sequence-specific elements in DNA. Only 3% of ER binding sites are in proximal promoter area (within 5 kb) of the targeted gene, while the rest are present at distal regulatory components. These are named enhancers (181, 182) and can regulate transcription processes over extensive distances of countless thousand base pairs and from a location downstream or upstream of the location of transcription initiation (207). Enhancer activity models have elucidated the looping of chromatin. An enhancer far away in the distance from its target promoter on a linear chromosome can be brought into proximity in 3D space (208).

Some studies established ER functions via chromatin looping that brings transcriptional regulatory complexes of enhancers with the specific target promoters, connecting the ERs activity at distal locations to transcriptional start sites (TSSs) (209, 210). Chromosome conformation capture (HiC) detects chromatin interactions and, with the use of deep sequencing, maps these interactions in the 3D organization of the genome. This recent advancement has revealed that most of the enhancer-promoter interactions happen within large organizing components (between hundreds of kilobases) of the genome designated as topologically associating domains (TADs) (211). The utilization of the same technique shows that E2-ER $\alpha$  prompts global chromatin restructuring in MCF7 cells (212) and reorganizes open and highly transcribed chromatins, permitting genes to share the transcriptional complex and regulatory elements as reviewed in (201).

Genome-wide studies also suggested that many enhancers and promoters contain a typical histone mark. Histones flanking active promoters are marked by H3K4me3, while enhancers are marked by H3K4me1, with both marked by H3K27ac after activation (213, 214). Moreover, within the locality of highly expressed genes, multiple transcription factors are recruited to initiate transcriptionally more active clusters of enhancers in relatively small sections of DNA. These are called super-enhancers which regulate cell type-specific and disease-related genes, including oncogenes (215, 216). Super-enhancers can be categorized as coregulated loci, by computationally detecting peaks containing high transcription factor binding levels and intersecting with dynamic chromatin marks, such as H3K27ac and DNase I hypersensitivity mark (215, 217).



Enhancer RNAs are also hallmarks of active enhancers and are involved in enhancer activation and functional activity (218, 219). These are non-coding RNAs that are transcribed at enhancers as a result of RNA polymerase II action. A fascinating likelihood is that enhancer RNAs might partake in a structural task in developing or preserving enhancer-promoter loops. Indeed, new data supports this theory (220, 221).

## **1.5 ER $\beta$ role in CRC**

### **1.5.1 Major challenges in the field**

It has been 25 years since ER $\beta$  was discovered; however, ER $\beta$  expression and distribution in tissues and cells is unclear and still debatable. ER $\beta$  expression and its role in MCF7 (breast) and LNCaP (prostate) cancer have been extensively studied, in contradiction with clinical parameters. Further, ER $\beta$  is not present at the mRNA level in most cell lines. Fifteen out of 934 cell lines show ER $\beta$  expression based on RNA-Seq data (222). Only B-cell lymphoma and ovarian granulosa cell lines expressed ER $\beta$ . Although ER $\beta$  antibodies detecting protein expression, irrespective of mRNA raises the question of antibody specificity. Nonspecific antibodies are indeed a major problem in research (223, 224).

Our group made efforts to study and validate extensively used ER $\beta$  antibodies. We reported that the most widely used ER $\beta$  antibodies (14C8, PPG5/10) are not specific to ER $\beta$  protein. We have identified that only one of 13 antibodies (PPZ0506) specifically targets ER $\beta$ , and this antibody was not widely used. Using this validated antibody (PPZ0506), we have detected ER $\beta$  expression in human testis, ovary, granulosa cell tumors, and colon (225, 226). During the same period, Nelson and colleagues also reported the use of unspecific ER $\beta$  antibodies in research (227). A comparative study between WB and IHC has shown that antibody performance depends on the application and suggested validating antibody for the intended application.

The use of unspecific antibodies in literature raises the question of functional significance of ER $\beta$ , and caution must be exercised in interpreting data obtained from such experiments. Therefore, unspecific antibody-based data was excluded from the literature review, and only results also supported by mRNA evidence were considered.

### **1.5.2 Epidemiological evidence**

It is a well-established fact that sex plays a vital role in CRC progression and survival. Men have increased progression and decreased survival compared to women of all ages (228, 229). Men with inflammatory bowel disease are at a 60% higher risk of developing colon cancer than women (20). Women benefit from significantly longer survival after rectal cancer resection than men. These results indicate that sex steroids may protect women from colorectal cancers (230, 231). Though not initially designed to explore CRC risk, the Women Health Initiative study established that use of hormone replacement therapy (HRT) protects women from CRC by reducing the risk up to 30% (232, 233). HRT is a combination of estrogen and progestin, and several studies have shown that estrogen alone contributes to large protective effects (234, 235). Also, the use of oral contraception has been shown to reduce CRC incidence by

approximately 14% (236). Consequently, oophorectomy (the removal of ovaries) in women increased CRC risk by 30% (237). These findings suggest that estrogen may protect against colon cancer, but the mechanisms by which this occurs remain unclear.

A meta-analysis study showed that soy consumption, which is rich in phytoestrogens, is associated with a 21% reduction of CRC risk in females but not in men, concluding that soy supplementation is synergistic with estrogen *in vivo* (238). Additionally, when individuals travel from an Asian nation to the United States, the occurrence of CRC in the immigrants increases to the average levels seen in the United States, indicating lifestyle and environmental factors role in etiology (239). Amongst the aspects related to the so-called *western* way of life, obesity occupies an important risk element regarding CRC. Notably, a positive relationship between the BMI and the frequency of colonic adenoma and advanced polyps was revealed in moderately young persons of both sexes and in premenopausal women with estrogen effects (240).

### 1.5.3 CRC suppression by ER $\beta$

While ER $\alpha$  is upregulated in several cancers, most notably in breast cancer, where it promotes cell proliferation, ER $\beta$  expression appears to decrease during cancer progression. At the mRNA level, ER $\beta$  is lost in CRC while expressed in non-tumor colon tissue. ER $\beta$  has been identified as the predominant ER type in colon in many studies, while ER $\alpha$  protein is not detected in the colonic epithelium or CRC (241).

In ER $\alpha$ -knockout mice using the AOM tumorigenesis model, soy protein was shown to protect ER $\alpha$ -knockout mice and WT mice from colon cancer (242). However, studies in ovariectomized Apc<sup>Min/+</sup> mice treated with coumestrol and genistein showed that coumestrol lowered tumor number in the colon and small intestine while genistein was found to have no significant effect (243). However, estrogen has been shown to mediate protective effects in wild-type mice, but not in ER $\beta$ -knockout mice, supporting ER $\beta$  as a mediator of these effects (244). Therefore, the protective effects of CRC from E2 could depend on colonic ER $\beta$  (245).

Saleiro et al. demonstrated that in females, whole-body deletion of ER $\beta$  causes an increase in colon adenomas and size in the AOM-DSS mouse model. Besides, they found that the loss of ER $\beta$  increased the levels of  $\beta$ -catenin and inflammation related molecules (246). However, using the same model, a study by Heiman et al. found that ovariectomy protected female mice from colitis-associated tumor development and estradiol and medroxyprogesterone acetate (MPA) or both promoted tumorigenesis. Using ER mutant mice, they concluded that estrogen depends on ER $\alpha$  and ER $\beta$  for protumorigenic effects (247).

A recent study using different colitis models induced by 2,4,6-trinitrobenzenesulfonic acid (TNBS) showed that E2 reduced the expression of pro-inflammatory cytokines. Interestingly, the authors also noted increased proliferation and reduced apoptosis in ER $\beta$ KO mice and suggested the differential protective effects of E2 in presence and absence of ER $\beta$  (248). Interestingly, ER $\beta$ KO mice exhibited increased cellular proliferation of colonic crypt cells in

both colitis models (DSS and TNBS), indicating the importance of ER $\beta$  in the protection against colorectal tumors (246, 248). Similarly, enhanced intestinal tumorigenesis was demonstrated in ER $\beta$  knockout ovariectomized Apc<sup>Min/+</sup> mice (249).

Along with those above epidemiological and animal studies about estrogen and ER $\beta$  protective effects, there is significant evidence supporting tumor suppressive activity of ER $\beta$  in cell line models. Re-introduction of ER $\beta$  in human colon cancer cell lines (SW480, HT29 and HCT116) downregulates IL6 inflammatory marker and reduces the expression of numerous oncogenes, for example, MYC, MYB and PROX1 (250, 251). Strikingly, ER $\beta$  has been shown to increase p53 signaling and reduce  $\beta$ -catenin levels, which increased apoptosis and reduced cell proliferation in LoVo colon cancer cells (252). These studies combined strongly support a role for intestinal ER $\beta$  as a target for chemo-preventive therapy in colon cancers, however, the molecular mechanisms behind ER $\beta$  function are not well understood.

#### 1.5.4 ER cross-talk

To coordinately regulate target genes, ER's recruit several transcription factors as partners to activate or repress hundreds of target genes at once (181, 190, 253). Many studies have shown a potential cross-talk between ER $\alpha$  and other transcription factors; only a few studies were conducted on ER $\beta$  cross-talk. Although ER $\beta$  shares several transcriptional mechanisms with ER $\alpha$ , there are unique ER $\beta$  mechanisms. ER $\beta$  undergoes conformational changes specific to each ERE, which result in the recruitment of differential coactivators (254). While both ERs have almost identical DBDs, ER $\beta$  has a lower affinity to ERE half-sites than ER $\alpha$  (255, 256). These differences between half-site binding and ERE-dependent conformational changes could contribute to the differences between ER $\alpha$  and ER $\beta$  in transcription activity (257).

Previous studies have identified many pioneering factors and coregulators crucial for transmitting the hormonal signal to the transcriptional machinery. FOXA1 is identified as a pioneer factor that binds to condensed chromatin allowing the recruitment of other transcription factors to the DNA (258). Particularly, FOXA1 reprograms ER $\alpha$  recruitment at *cis*-regulatory components in breast cancer cells (182). Numerous sequencing initiatives, along with further research, identified regulators of ER $\alpha$  (FOXO3A, FOXM1, GATA3, ERBF-1) (259), as well as ER $\beta$  (CREB in endometrial cells (260), AP2 $\alpha$  in prostate cancer cells (261), and Clock/*Bmal1* in mouse models (262). Recently a couple of studies reported that tet methylcytosine dioxygenase 2 gets involved in ER complex along with GATA3 and acts as a necessary component for ER binding to chromatin (263, 264).

Several studies reported that different steroid receptors and transcription factors modulated p65 chromatin binding. In the breast cancer field, numerous studies indicated that ER $\alpha$  represses p65 to exert anti-inflammatory activity (265-268), whereas some studies have shown that ER $\alpha$  and p65 work together to increase gene transcription synergistically (189, 269-274). It is, therefore, unclear whether ER $\alpha$  and p65 activate or repress each other, but an interaction between ER $\alpha$  and p65 has been consistently reported. In a study, Franco et al. (275) revealed that TNF $\alpha$ -activated NF $\kappa$ B reprograms ER $\alpha$  to a subcategory of NF $\kappa$ B-binding sites, resulting

in increased production of enhancer RNA transcripts, an activity linked beforehand to enhancer activation by ER (276).

The interaction between ER $\alpha$  and NF $\kappa$ B is shown at the baculoviral inhibitor of apoptosis 856repeat-containing 3 (BIRC3) target gene. TNF $\alpha$  activated NF $\kappa$ B binds to the BIRC3 promoter and prepares it for ER $\alpha$  binding, which in return enhances TNF $\alpha$ -induced BIRC3 activation (277). Besides, TNF $\alpha$  intensely transforms the ER $\alpha$  enhancer setting in an NF $\kappa$ B-dependent fashion (275, 278). Our group has reported that also ER $\beta$  regulates NF $\kappa$ B target genes, including downregulation of Il-6, which is the main target of the NF $\kappa$ B pathway (279). However, there is no data available for ER $\beta$  coactivators in colon cells, thus understanding the cross-talk between ER $\beta$  and its co-activators is essential for understanding its function. Based on that, in this thesis, I am exploring whether ER $\beta$  and NF $\kappa$ B interact specifically in colon cancer. Learning more about NF $\kappa$ B signaling and how it is influenced by estrogen signaling should lead to better therapeutic approaches in the future to target it in specific cancer types.

### 1.5.5 ER cistrome studies

Albeit ER $\alpha$  inhibitors have been utilized in clinics since the 1980s, the mechanism of ER $\alpha$ 's genomic activity on a genome-wide scale remained mysterious for a long time (280). Primary progress of chromatin immunoprecipitation (ChIP) together with DNA microarray analysis (ChIP-on-chip) techniques, ER $\alpha$  genomic binding, was identified on a genome-wide scale for the first time (182). Through the expansion of high-throughput sequencing methods, several ER $\alpha$  ChIP-Seq studies were executed in cell lines and some in tissues (281). Almost 95% of the entire ER $\alpha$ -binding locations are premised at distal *cis*-regulatory components (henceforth labeled as 'cistromics') that later acquired recognition as enhancer areas (280, 282).

Several studies have been published on ER $\beta$  ChIP-Seq in breast mainly using exogenous expression of ER $\beta$ . In MCF7 cells, studies using different antibodies showed between ~400 to 38 000 ER $\beta$  binding sites (235, 283-285). Several studies in different cell lines (286) or tissues (287, 288) identified ER $\beta$  binding sites somewhere between 5000 to 14 700 binding sites; few studies used antibodies not highly specific to ER $\beta$ . Though differing studies indicate changeable statistics of ER binding actions (even in the same cell line), possible because numerous enhancers might control a single gene (289), or certain variances are attributable to the experimental conditions and the lack of specificity of antibodies that were utilized.

## **2 RESEARCH AIMS**

### **General aim**

In epidemiological and animal studies, estrogen receptor beta (ER $\beta$ ) has been shown to protect from colorectal cancer (CRC), but the mechanisms are poorly understood. Thus, the overall aim of this thesis is to elucidate the role of ER $\beta$  and evaluate its potential as a therapeutic target in prevention of colorectal cancer. The research articles presented in this thesis identify molecular mechanisms of ER $\beta$  involved in preventing CRC formation and development and propose novel insights involved in CRC prevention.

### **Specific aims**

Paper I: To test if intestinal epithelial ER $\beta$  mediates protective role in CRC, using a colitis-induced tumor mouse model with tissue specific knockout of ER $\beta$  and to investigate any sex differences.

Paper II: Determine the genome-wide binding pattern and regulation of ER $\beta$  in colon cancer cells and identify its cofactors.

Paper III: Elucidate the impact of estrogen signaling in HFD induced colon inflammation in both sexes.

Paper IV: Evaluate the cross-talk between ER $\beta$  and p65/TNF $\alpha$  inflammatory signaling in colon cancer cells.

### 3 MATERIALS AND METHODS

#### 3.1 Mouse models

Fundamentally, animal models of human disease are valuable models for studying several diseases, including the development of cancer. Animal models also play a crucial role in translating scientific studies into human disease and in predicting the future of novel therapeutic drug targets. The animal studies got approved by the Swedish Regional Board of Animal Research. In Sweden, the use of laboratory animals requires well-controlled ethical permission, controlled premises, and adherence to high animal welfare requirements. In the use of animals, the "3Rs" recommendations that encompass Replacement, Reduction, and Refinement were adopted in 1959 (290). Replacement: Animal experimentation is avoided wherever possible, and *in vitro* models should be used instead. Reduction: use the smallest number of animals required for statistical power. Refinement: reduce animal suffering and improve animal welfare.

The mice were constantly provided with a regular chow diet and water and were kept on a 14-hour light, 10-hour dark cycle. Primarily, in the paper I, CA-CRC was induced in WT mice and mice that specifically lack ER $\beta$  in intestinal epithelial cells (referred to as ER $\beta$ KO<sup>Vil</sup>). Regarding paper III, C57BL/6J mice with HFD-induced obesity were used.

##### 3.1.1 ER $\beta$ KO<sup>Vil</sup> mouse

ER $\beta$ KO<sup>Vil</sup> mouse model maintained on C57BL/6J background strain was generated by Cre/loxP recombination technology. ER $\beta$ flox/flox mice with ER $\beta$  exon 3 (B6.129X1-Esr2<sup>tm1Gust</sup>) flanked by two loxP sites were bred with transgenic mice expressing Cre recombinase under the control of the enterocyte-specific Villin promoter. Cre is therefore only expressed in the intestine, where it deletes the DNA sequence between the loxP sites in a site-specific manner. ER $\beta$  gene exon 3 generates the first zinc finger in the DBD. Its knockout led to frameshift in the coding region following splicing from exon 2 to exon 4, resulting in a frameshift and an early stop codon, reduced mRNA levels (due to nonsense-mediated decay), and no ER $\beta$  protein expression. ER $\beta$ flox/flox mice without Cre allele were used as controls (denoted as WT). The opportunity to characterize the function of ER $\beta$  specifically in the intestinal epithelium is a benefit of ER $\beta$ KO<sup>Vil</sup> mice. It is also worth noting that ER $\beta$  is lost in the small intestine. The full-body knockout of ER $\beta$  in mice results in phenotypes with fertility issues with few littermates (291), but so far, no major effects were observed in the colon of ER $\beta$  knockout mice in standard conditions. However, challenging ER $\beta$  knockout mice with inflammatory conditions or obesity mice gives more pronounced effects due to lack of ER $\beta$ .

##### 3.1.2 AOM/DSS-induced colitis

Azoxymethane (AOM)/dextran sodium sulfate (DSS) model has been used extensively in research to mimic human CA-CRC features. The AOM/DSS induces distally located invasive tumors and molecular modifications in  $\beta$ -catenin, pro-inflammatory cytokines (TNF $\alpha$ ), similar to human CA-CRC (292-294). ER $\beta$ KO<sup>Vil</sup> and WT mice between 5 and 10-weeks were

intraperitoneally injected with colonic carcinogen AOM (10mg/kg) on day 1 and then accompanied by colonic irritant DSS treatment in drinking water starting from day 7 to 14. The DSS cycle was then repeated three times, with two weeks of regular drinking water in between. For the duration of the DSS treatment, the mice lose weight, have diarrhea, and develop rectal bleeding, and all these different parameters were monitored using a score sheet. The mice recover during the phase of drinking normal water and, in due course, acquire colitis followed by adenomas. Following 9 or 15-16 weeks of AOM injection, the mice were sacrificed. Notably, the AOM/DSS technique has excellent reproducibility and potency, besides being a simple and inexpensive method of application. It is worth noting that different strains of mice respond differently to this treatment and that their microbiota composition can influence the efficacy of tumor development.

### **3.1.3 High-fat diet (HFD) induced colon inflammation and obesity**

Male and female C57BL/6J mice bred in-house at the animal facility and aged 5 to 6 weeks were fed an HFD (60% kcal fat) or a control diet (CD) (10% kcal fat) for 13 consecutive weeks. The mice were treated with vehicle or estrogenic ligands, E2 (0.05 or 0.5 mg/kg), PPT (2.5 mg/kg), and DPN (5 mg/kg) for a total of 9 injections for 3 weeks before sacrifice. The ligands were prepared in a mixture containing 40% PEG400, 5% DMSO, and 55% water. In comparison to the regular western diet, which has approximately 45% calories from fat, this diet has a higher fat content. The physical properties of the HFD diet, which easily breaks into powder, creates a problematic situation regarding monitoring the food intake. Measuring their exact caloric intake, and controlling their actual consumption, is therefore, a considerable challenge (295).

### **3.1.4 Tissue collection**

Weighed colons were cleaned and opened along the axis, and the number and size of adenomas were recorded. The colons were fixed with 4% formaldehyde for 24 hours and kept in 70% ethanol before embedded in paraffin for immunohistochemistry (IHC) or the fresh tissue is directly frozen in liquid nitrogen for analysis, including with RNA extraction and quantitative reverse-transcription PCR (qPCR). Also, fresh tissues were directly embedded in OCT and frozen on dry ice and stored at 80°C. A pathologist who was unaware of the mouse genotype scored the colitis in hematoxylin and eosin (H&E) stained sections. The colitis score (0-12) was calculated: the severity, the extent of hyperplasia, level of ulceration, and the size of the affected areas were all graded between 0-3.

## **3.2 *In vitro* studies**

### **3.2.1 Colon cancer cell lines**

Well-characterized cell lines enable reproducibility and resolve most of the ethical issues that come with using animal and human tissues. Due to these reasons, cell line models continue to be the most widely used in research worldwide. The foremost objective in choosing CRC cell lines HT29 and SW480 in Paper II and IV was to create an optimized, well-controlled system

for studying the genome-wide binding of ER $\beta$ . HT29 is an epithelial-like human colorectal adenocarcinoma cell line generated from a 44-year-old female colon adenocarcinoma tumor (296). The SW480 was obtained from a 50-year-old male with type B colorectal adenocarcinoma (297). Differences include that HT29 cells are CIMP (CpG island methylator phenotype) positive, whereas SW480 cells are CIMP negative. The HT29 contains WT KRAS, PTEN genes, and mutations in BRAF (V600E), PIK3CA (P449T), and TP53 (R273H). SW480, on the other hand, has WT BRAF, PIK3CA, and PTEN and mutations in KRAS (G12V) and of both TP53 alleles (R273H and P309S) (298). The P53 allele in the SW480 cell line is a double mutant, while the P53 allele in the HT29 cell line is a single mutant, which may result in a different chromatin complex with P65 (298). We also used ER $\alpha$ -positive MCF7 breast cancer cells with tetracycline-induced expression of ER $\beta$  in paper II to compare tissue-specific ER $\beta$  chromatin-binding between colon and breast. In Paper IV, we used MCF7 parental cells to compare p65 cistrome between colon and breast. Due to the lack of ER $\beta$  in CRC cells, ER $\beta$  has been exogenously added using a lentivirus system, which has been generated and characterized previously (250, 251, 299).

Cell lines were cultured in a standard cell culture medium comprising 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin. The cells were serum-starved using 1% charcoal-stripped FBS and DMEM-phenol red-free media. Regular FBS contains hormones, such as estrogen and phenol (which have weak estrogenic properties). *In vitro* models are an excellent choice to dissect molecular mechanisms; however, they have a drawback when studying a complex disease like CRC because of the lack of tissue microenvironment that influences tumor growth and metastasis *in vivo*.

### 3.2.2 Chromatin immunoprecipitation

In many major cellular processes, such as transcription, splicing, replication, and DNA repair, DNA-binding proteins play crucial roles. These proteins include transcription factors that preferentially bind to specific sequences of DNA and histone proteins that form the nucleosome core, which is the chromatin's basic unit. To study genome-wide binding of transcription factors, chromatin immunoprecipitation (ChIP) combined with microarrays (ChIP-on-chip) or sequencing (ChIP-Seq) has become standard techniques (300, 301). In brief, the cells were cross-linked to capture protein-DNA interactions, and the chromatin was sheared into small fragments in the range of 200-600 bp by sonication. DNA-protein complexes were immunoprecipitated with an antibody specific to the protein of interest. Cross-links were reversed, and the released DNA was sequenced to determine the sequences bound by the protein (301, 302). ChIP-Seq performed with a specific antibody provides an accurate map of binding sites for transcription factors and other DNA-binding proteins, and this information is crucial for understanding biological processes. Integrating ChIP-Seq data with functional genomic and transcriptomic data (RNA-Seq) can provide a global view of chromatin accessibility and regulation of gene expression (303). ChIP-Seq is the primary technique used in papers II and IV, where it is accompanied by RNA-Seq or microarray data.



It is important to remember that each method has its limitations. From a technical perspective, for a high-quality ChIP experiment, a highly specific antibody and a large number of cells (~10 million) are usually required (304, 305). In addition to direct binding events, the signal from ChIP-Seq also includes indirect binding in which one factor interacts with another DNA-binding factor (tethering). It is essential to distinguish between direct DNA binding and tethering events, but this is not directly achieved from ChIP-Seq data but requires further analysis, *in silico* or experimental (301). Recent technological advances in the ChIP-Seq protocol have made it possible to test samples with limited cells, increase the accuracy of the genomic location of binding events, and test multiple binding events (306-309).

The ChIP-Seq data pipeline addresses various aspects of the assessment and management of data: the identification of artifacts, filtering low-quality reads, estimating library complexity, sophisticated reading alignment and peak calling algorithms, and downstream analytical tooling (300). For a ChIP-Seq comparison, a common normalization approach uses the total number of mapped reads per million to adjust for biases generated by sequencing depth between samples. This approach works only to correct technical artifacts but cannot correct biological differences like different peak numbers. Several publications have pointed out these biases (310-313) and indicated the need for an initial normalization step. Various strategies have been employed, such as relative level difference and absolute level difference using synthetic histone spike in normalization (314, 315). Recently, another normalization strategy called parallel factor ChIP addressed these challenges by using a second antibody against the target chromatin as an internal control (316). Although the need for an exogenous spike-in is eliminated in this technique, its dependence on antibodies might cause reproducibility concerns. There are no gold standards available for ChIP-Seq data normalization and analysis. Hence the optimal method should be selected based on prior system knowledge and sequenced sample statistics (300, 302). There are web-based services such as ChIPseeker (317), Nebula (318), and CSA (319) that are available to analyze ChIP-Seq data.

The first method for collecting single-cell chromatin data has been published using the "Drop - ChIP" method (320). This method uses a droplet microfluidic labeling system for individual cell chromatin before immunoprecipitation and reports ~800 reads per cell. Recent development using the microfluidic droplet method has a higher resolution with ~10,000 reads per cell. Single-cell ChIP-Seq using this method was performed in breast cancer and provided a new alternative to study complex chromatin heterogeneity in tumors (321).

### **3.2.3 Luciferase assays**

We performed luciferase assays to determine how ER $\beta$ , AP1, or NF $\kappa$ B can affect transcription and potentially cross-talk. The reporter gene expression is kept under the control of firefly or renilla luciferase. We used luciferase constructs with estrogen response element (ERE), NF $\kappa$ B response element (NF $\kappa$ B-RE) (Paper I), or TPA response element (TRE) (Paper II) in their

promoters. Plasmid with renilla luciferase reporter was used as a control to normalize enzyme activity of firefly luciferase.

Firefly and renilla luciferase are two bioluminescent forms of proteins that emit light in response with their substrate D-luciferin (firefly) or coelenterazine (renilla), and the produced light intensity is proportional to the protein concentration in the sample. The reporter gene's expression is directly proportional to the protein expression allowing for monitoring of small changes in expression.

#### **3.2.4 Statistical analysis**

For statistical analysis, GraphPad Prism was used (GraphPad Software Inc). The data is provided as a mean  $\pm$  standard error of the mean (SEM). For comparison between two groups, a two-tailed Welch's t-test or paired t-test (if paired data) was used. For comparisons between multiple groups, one-way and two-way analysis of variance (ANOVA) were used, followed by Fisher's LSD test for *in vivo* data or Tukey's multiple comparisons test.

## 4 RESULTS

### 4.1 Paper I: Intestinal estrogen receptor beta suppresses colon inflammation and tumorigenesis in both sexes


Studies in full-body ER $\beta$  knockout mice and selective activation of ER $\beta$  with ligands have previously confirmed the notion that estrogen via ER $\beta$  protects against CRC (244, 322-325). Our group's research with exogenously expressed ER $\beta$  has shown the antiproliferative and antitumorigenic effects of ER $\beta$  in both colon cancer cell lines (326-329) and xenografts (327). Thus, selective activation of ER $\beta$  is a potential choice for CRC preventative therapy, but its mechanism of action needs to be elucidated. ER $\beta$  has been detected in a few immune-related organs and some immune cells (225), and it is possible that ER $\beta$  could impact CRC development through the immune system. The previous studies which explored the role of ER $\beta$  in CRC using full knockout animals could not exclude possible systemic ER $\beta$  effects via the immune system. Also, with one exception (324), all studies were conducted only in female mice (244, 325). However, ER $\beta$  is expressed in colon and rectum in both men and women, men also synthesize estrogen (primarily E1), and ER $\beta$  can be activated by dietary phytoestrogens. Thus, men are also expected to have some ER $\beta$  activity. Hence, we here attempt to understand the physiological relevancy of ER $\beta$  in both sexes using colitis-induced CRC (CA-CRC) in male and female mice that lack ER $\beta$ , specifically in intestinal epithelium (ER $\beta$ KO<sup>Vil</sup>). We have used the AOM/DSS model to induce CA-CRC, which reflects the histopathological and molecular environment of human CA-CRC.

Using RNAscope and IHC with a validated antibody (225), we confirmed ER $\beta$  expression in normal colon and rectum epithelia of both human and mouse, and lack of expression in CRC, which supports our underlying hypothesis that ER $\beta$  can protect against CRC formation. We observed that loss of intestinal ER $\beta$  increases the number of tumors in males and the tumor size in females (Figure 6). One interesting finding is that loss of ER $\beta$  increased the pro-inflammatory cytokine *Tnf $\alpha$*  expression in both sexes, most prominently in males. Moreover, we noted that several TNF $\alpha$  and NF $\kappa$ B target genes, such as *Il6* and *Il1b*, strongly increased upon loss of ER $\beta$ , following AOM-DSS treatment, especially in males. On the other hand, in females, lack of epithelial ER $\beta$  increased both ulcerative areas and reduced ulcer healing. We have thus demonstrated that both sexes are protected by intestinal ER $\beta$  from CRC in a sex-dependent manner. The tumor size and incidence differences between males and females may be partly due to these specificities or microenvironmental signals, and differences in their immune system, resulting in different colon epithelial gene expressions between males and females.

We have also previously shown that the microbiota diversity in these mice appeared modulated by ER $\beta$  (330). To circumvent possible impact by microbiota effects and explore if ER $\beta$  could protect the epithelial cells from TNF $\alpha$ -induced damage, we treated intestinal organoid cultures generated from WT and ER $\beta$ KO<sup>Vil</sup> mice of both sexes, with TNF $\alpha$  and a selective agonist for ER $\beta$  (DPN). The *ex vivo* crypt formation, which quantifies regenerative growth, was decreased by TNF $\alpha$  treatment in both sexes of WT and ER $\beta$ KO<sup>Vil</sup> mice. In both sexes of WT mice, ER $\beta$ -

selective agonist DPN significantly counteracted the TNF $\alpha$ -induced epithelial cell damage but did not do so in ER $\beta$ KO<sup>Vil</sup> mice. This data demonstrates that the activation of intestinal ER $\beta$  protects both sexes against TNF $\alpha$ -induced epithelial cell damage.

Further, we explored these molecular mechanisms in two human colon cancer cell lines HT29 and SW480, using transcriptomic and genome-wide studies. We found that ER $\beta$  modulated a proportion of the TNF $\alpha$ -regulated transcriptome and inhibited most of these genes in CRC cells. Next, we tested whether ER $\beta$  can impact the translocation of NF $\kappa$ B subunit p65 or transactivation of TNF $\alpha$ . We observed that ER $\beta$  affected neither the translocation of p65 nor the transactivation of TNF $\alpha$  in either of the cell lines. We could confirm that ER $\beta$  modifies the expression of NF $\kappa$ B regulated genes also in mice, such as *Atf3*, *Bcl3*, and reduces pro-inflammatory signaling by lowering levels of chemoattractants *Ccl2* and *Ccl4*. NF $\kappa$ B targets ATF3, BCL3, and BIRC3 also had ERE elements in close proximity, indicating that ER $\beta$  has the potential to bind these targets. Using ChIP-qPCR, we have confirmed that ER $\beta$  binds to NF $\kappa$ B targets, ATF3, BCL3, and BIRC3 in the promoter or enhancer regions, indicating biological cross-talk between ER $\beta$  and p65. Further, using the ERE-luciferase transactivation assay in cell lines, we found that TNF $\alpha$  could enhance the transactivation of ER $\beta$ . This suggests that an inflammatory condition increases ER $\beta$  activity, which successively inhibits TNF $\alpha$ -mediated signaling and thus reduces the inflammation, implying an inhibitory feedback loop.



**Figure 6.** (A, B) Intestinal ER $\beta$  protects both sexes from colitis-induced CRC in a sex-dependent manner. Loss of intestinal ER $\beta$  increases the number of tumors in males while increasing the tumor size in females (226).

In this study, we tested only a limited number of promoter regions. Further performing ER $\beta$  and p65 ChIP-Seq is significantly more informative and reveals the molecular basis for this cross-talk, which is addressed in paper II and IV. In conclusion, our data indicate that ER $\beta$ , expressed specifically in the intestinal epithelia, protects from colitis-induced CRC in a sex-dependent manner by attenuating TNF $\alpha$ - NF $\kappa$ B pathway.

## 4.2 Paper II: Genome-wide estrogen receptor $\beta$ chromatin binding in human colon cancer cells reveals its tumor suppressor activity

The antitumorigenic effects of ER $\beta$  have been previously reported by our group in different CRC cell line models, where re-expression of ER $\beta$  reduced cell proliferation, migration, and invasion (245, 250, 251, 299). In paper I, the deletion of intestinal ER $\beta$  in mice enhanced tumorigenesis, indicating the importance of colonic ER $\beta$  in protecting from tumors. All these studies strongly support a role for intestinal ER $\beta$  as a target for chemopreventive therapy in colon cancers; however, the ER $\beta$  mechanisms of action at the chromatin level in colon cells have not been studied yet. So far, only a few studies have mapped ER $\beta$  binding sites, mostly in MCF7 breast cancer cells (16, 17). The ER $\beta$  chromatin binding in the colon is unknown.











To determine colonic tumor protective effects of ER $\beta$  through its activity as a transcription factor, we aimed to characterize its genome-wide chromatin binding sites and identify its target genes in colon. One of the challenges in studying ER $\beta$  is its low native expression in tissues and its lack of expression in immortalized cancer cell lines (257). Introducing ER $\beta$  into cell lines is a way of restoring its loss of endogenous expression to study its impacts. For this purpose, we performed ChIP-Seq on previously characterized CRC cell lines with transduced ER $\beta$  expression, HT29-ER $\beta$ , and SW480-ER $\beta$  (299, 327). The quality of a ChIP-Seq study is highly dependent on the specificity and selectivity of the antibody used, and we further optimized and evaluated the highly validated ER $\beta$  antibody PPZO506 for application in ChIP-Seq (225).

We mapped 2977 and 1853 ER $\beta$  binding sites in HT29, SW480 cells, respectively. In line with previous ER cistrome studies, most ER $\beta$  binding sites were located within intronic (43%) and intergenic (40%) regions. We found that the majority of binding sites contain canonical ERE motif, indicating that ER $\beta$  binds directly to DNA with its DNA-binding domain. Further, a relatively large proportion exhibited AP-1 motif, indicating its known interaction with the AP-1 complex. We also discovered motifs (TCF, ELF3, and KLF5) that have not previously been associated with ER $\beta$ , indicating potential novel cross-talk (Figure 7). Overall, the most enriched biological functions among genes closest to ER $\beta$  binding sites include cell migration, cell-cell adhesion, and transcriptional regulation in both cell lines. These are processes that ER $\beta$  is known to affect.

To determine whether ER $\beta$  binding is functionally relevant in transcriptional regulation of genes, we linked ER $\beta$  ChIP-Seq data with corresponding gene expression data from SW480 and HT29 cells. Associated with ER $\beta$  binding sites, ER $\beta$  regulated genes are mainly involved in proliferative Wnt signaling, NF $\kappa$ B signaling, and insulin-like growth factor signaling pathways. We identified that ER $\beta$  strongly upregulated CST5, a candidate tumor suppressor in colon cancer, and downregulated the LRP6 oncogene, which promotes aberrant  $\beta$ -catenin signaling. To transcriptionally regulate the oncogenic Wnt pathway,  $\beta$ -catenin depends on TCF family proteins, as it lacks a DNA-binding domain (331). A significant fraction of ER $\beta$ -binding sites contained TCF4 motif in both SW480 (12%) and HT29 (19%) cells.

To investigate whether ER $\beta$  impacts the epigenome, we studied the histone acetylation mark H3K27ac which represents transcriptionally active enhancers, promoters, in presence and absence of ER $\beta$ . A significant fraction of H3K27ac binding sites were modulated by the presence of ER $\beta$  in both HT29 (4159 enhanced, 4835 reduced) and SW480 cells (3793 enhanced, 3596 reduced). Genes located by ER $\beta$ -modulated H3K27ac sites were also overrepresented among functions such as cell migration, apoptosis, and Wnt signaling.

To explore whether or to which extent ER $\beta$  binding is tissue-specific, we compared our ER $\beta$  cistrome data with that of breast cancer cell line MCF7, using the same antibody and method. The predominant ER $\beta$ -binding motifs in breast were ERE, AP-1, and GATA4. We found no enrichment for TCF and KLF motifs in breast, indicating that ER $\beta$  tethering with TCF and KLF5 is specific to colon cancer and thus cell context dependent. TCF and KLF5 chromatin accessibility was dysregulated in polyps associated with CRC invasion, according to a recent study that used ATAC-Seq to construct single-cell chromatin maps of polyps and CRC. TCF motifs were highly enriched in polyps, indicating increased chromatin accessibility, whereas KLF motifs chromatin accessibility was lost during cancer progression (332). KLF4 was also identified as a potential tumor suppressor in CRC (333). One could speculate that ER $\beta$  directly or indirectly competes with  $\beta$ -catenin to bind TCF proteins and controls the genes or regulatory elements involved in the early events of polyp transforming into CRC.

HT29 ER $\beta$ ChIP	Motif	P Value	% sequences	SW480 ER $\beta$ ChIP	Motif	P Value	% sequences
	ERE(NR)	1e-950	77%		ERE(NR)	1e-984	55%
	AP-1	1e-415	28%		TCF	1e-74	19%
	TCF	1e-76	34%		RUNX2	1e-70	18%
	ELF3	1e-59	20%		ELF3	1e-23	20%
	KLF5	1e-50	24%		AP1	1e-21	10%

**Figure 7.** *De novo* motif analysis identified TCF and AP-1 motifs in ER $\beta$  binding sites in both colon cancer cell lines (334).


The use of exogenously expressed ER $\beta$  instead of native expression is one of the shortcomings of this study, and also that the cell lines are from colon cancer instead of non-tumor colon epithelial origin. However, the levels of ER $\beta$  in normal epithelial and non-tumor cell lines are too low to allow high-quality ChIP-Seq data, and non-tumor colon cell lines are difficult to culture and transduce. In conclusion, this study mapped genome-wide ER $\beta$  binding sites and identified the epigenetic changes by ER $\beta$  in colon cancer cell lines for the first time.

#### 4.3 Paper III: High-fat diet and estrogen impact the colon and its transcriptome in a sex-dependent manner

Sex and lifestyle factors, like dietary factors, are important risk factors in CRC development. Intake of a High-fat diet (HFD) affects the colon first and modulates intestinal permeability, microbiota composition, and induces inflammation (81-83). A meta-analysis of 56 cohort and case-control studies found that obesity increases the risk of CRC, and a 5 kg/m<sup>2</sup> increase in body mass index (BMI) increases the risk of CRC development by 18%, and a BMI above 30 kg/m<sup>2</sup> increases the risk by 41% (79). Obesity impacts inflammation, gut microbiota, hormones such as estrogen, insulin, and leptin which all play a role in CRC (335, 336). Estrogen has shown to protect against obesity and metabolic syndromes, as reviewed in (337). In study I, we also found that ER $\beta$  protects from CRC in a sexually dimorphic manner, with more tumors in males and larger tumors in females. However, no studies have been conducted to investigate colonic sex differences in relation to obesity, metabolic dysregulation, and estrogenic ligands. In this study, we focused on determining if estrogen affects the colon microenvironment during HFD-induced inflammation and on dissecting sex differences. For 13 weeks, we fed mice an HFD with 60% fat or a control diet (CD) with 10% fat. In the last three weeks prior sacrifice, mice were treated with different estrogenic ligands. To investigate sex differences on the colon transcriptome, we performed RNA-Seq on male and female distal colon tissues.

The colon transcriptome of CD-fed mice revealed that a high proportion of genes (1564) were differentially expressed between males and females. Under HFD treatment, the sex difference was less prominent, with 364 differentially expressed genes. About a quarter of the transcriptomic sex differences found under HFD were also observed under CD. Among these stable sex difference genes, the most overrepresented genes belonged to the circadian rhythmic pathway and included key genes *Npas2* and *Bmal1(Arntl)*, which were higher expressed in females compared to males. Further, separate comparisons showed that HFD modulated different genes in the sexes. Significantly overrepresented functions of genes regulated in males were cell cycle and hypoxia, while for females, it was lipid metabolism, steroid hormone, and Wnt signaling. Hypoxia can induce inflammation, and in IBD, hypoxia-inducible factors (HIF-1 $\alpha$  and HIF-2 $\alpha$ ) are highly present (338). In paper I, we also found strong inflammatory signaling and significantly increased *Il6* and *Il1b* in WT males compared to WT females, suggesting sex differences. We also identified that HFD-induced sex differences in colon length (a shortened colon is indicative of inflammation), fasting glucose, and crypt proliferation. A shorter colon was found only in males but not in females. Further, HFD increased crypt proliferation and fasting glucose levels in males but not in females. On the contrary, HFD enhanced F4/80<sup>+</sup> macrophage infiltration in both sexes but more strongly in females.

To determine if ER $\beta$  could modulate inflammation, we treated mice with estradiol (E2, activates all ERs), a ligand selective for ER $\alpha$  (PPT), and a ligand selective for ER $\beta$  (DPN). Treatment with E2 opposed the HFD-induced gene expression, particularly on genes involved in metabolic processes and inflammation. These results clearly indicate that E2 opposes the HFD dysregulated transcriptome, possibly through ERs present in the colon or through improved metabolism in general (Figure 8A). E2 and PPT both controlled body weight gain and total fat, which impacts the overall metabolic profile, which is in accordance with previous studies indicating that E2 through ER $\alpha$  reduces food intake (339) and regulates the body fat distribution (340). The impact in the colon was mainly seen through E2 and DPN, which counteracted HFD-induced F4/80<sup>+</sup> colon macrophage infiltration in both sexes and Ki67 proliferation in colon crypts of males. All three ligands E2, DPN, and PPT significantly restored the HFD reduced *Bmal1* gene in females. We further identified ER $\beta$  binding to BMAL1 and NPAS2 using ChIP-Seq data (from paper II) and correspondingly decreased gene expression of *Bmal1* and *Npas2* in the female colon with loss of intestinal ER $\beta$  (Figure 8B-D). This study concludes that HFD impacts the colon microenvironment with prominent sex differences. Estrogen modulates the HFD induced transcriptome and, through ER $\beta$ , counteracts HFD-dysregulated clock genes, macrophage infiltration, and cell proliferation in a sex-dependent manner.



**Figure 8.** (A, B) E2 opposes the majority of HFD induced gene dysregulation in colon microenvironment, such as dysregulation of clock genes BMAL1 and NPAS2 in males (C). Interestingly, we identified ER $\beta$  binds to clock genes in colon cancer cells using ChIP-Seq (D) The clock genes were dysregulated specifically in the absence of intestinal ER $\beta$  upon HFD treatment, indicating direct regulation of core clock genes in females by ER $\beta$  (341).



#### 4.4 Paper IV: Estrogen receptor beta influences the inflammatory p65 cistrome in colon cancer cells.


In paper I, we showed that ER $\beta$  protects against epithelial damage by repressing TNF $\alpha$  signaling *in vivo*. ER $\beta$  also binds in the vicinity of NF $\kappa$ B target genes and modulates the expression of TNF $\alpha$  regulated transcriptome in CRC cell lines. In paper II, we also find that ER $\beta$  chromatin-binding sites were enriched adjacent to genes involved in NF $\kappa$ B signaling. The ERs can interact with other transcription factors to promote or suppress transcription. In particular, a cross-talk between ER $\alpha$  and NF $\kappa$ B in breast cancer has been identified, and some studies (191, 192) support that ER $\alpha$  inhibits NF $\kappa$ B activity; on the contrary, other studies claim that ER $\alpha$  promotes NF $\kappa$ B activity (193, 194). The DNA binding domain is highly conserved between ER $\alpha$  and ER $\beta$ , but the two proteins differ significantly in their N-terminal domains and their ligand-binding domains (134, 136). Therefore, we hypothesized that ER $\beta$  transcriptionally modulates NF $\kappa$ B signaling by modulating its chromatin binding at the genome-wide level, for example, through protein-protein interaction or chromatin-binding competition. To experimentally demonstrate interactive cross-talk between p65 and ER $\beta$ , we performed p65 ChIP-Seq in CRC cells upon TNF $\alpha$  treatment in presence and absence of ER $\beta$  and compared with corresponding gene expression data.

We identified 3151 and 1459 p65 binding sites in HT29 and SW480 cells. They were located by genes involved in known biological functions such as TNF $\alpha$  signaling, cell migration, apoptosis, and more particularly circadian rhythm. The circadian rhythm has not been linked to p65 before, and this novel potential role in the colon paves the way for new studies. To compare p65 cistrome between different tissues, we have used publicly available p65 ChIP-Seq data from MCF7 cells (also performed using double crosslinking). Comparison of p65 chromatin-binding sites revealed that to a large extent, they are different between the breast and colon cancer cell lines, indicating that NF $\kappa$ B works uniquely to control transcriptional regulation in different cancers. The two p65 ChIP-Seq studies used different antibodies and protocols, which might also impact this comparison.

Further, we identified that ER $\beta$  modulates p65 chromatin binding in both CRC cells but in a cell line-specific manner. In paper III, we showed that ER $\beta$  restores the core clock genes dysregulated by HFD. Here, in paper IV, we found that ER $\beta$  modulates the p65 binding to core circadian genes CLOCK and BMAL2, further strengthening the intrinsic cross-talk between ER $\beta$ , clock genes, and p65. In HT29 cells, ER $\beta$  generally reduced p65 chromatin binding while it increased p65 binding in SW480 cells. However, in terms of gene expression, ER $\beta$  reduced p65 transcriptional activity in both cell lines. It is possible that biological, genetic, or epigenetic changes between HT29 and SW480 cell lines impact this cross-talk. Some notable differences are that the SW480 cell line contains double mutant p53 alleles (298), whereas HT29 has a single p53 mutation, which might result in different chromatin complexes with p65. Previous studies showed p53 and p65 interactions in various cancers with different outcomes (342-344), and it was shown in SW480 cells the double mutated p53 variants were able to impact p65

binding (345). Also, HT29 is a female cell line, while SW480 has a male background. Our results should be verified with more cell lines to further understand this cross-talk.

Some methodological and potential shortcomings of this study include that cells were treated with different TNF $\alpha$  timings and concentrations for the ChIP-Seq study compared with the gene expression experiments. Genomic binding occurs immediately after p65 translocation, and we have therefore used 30 minutes TNF $\alpha$  treatment for ChIP-Seq experiments. The regulation of gene expression by p65 is a slightly later event as it occurs after chromatin binding and the mRNA remains in the cell for a longer time; hence we used a 2-hour treatment for gene expression analysis. In paper II, we demonstrated that ER $\beta$  chromatin binding occurs independently of ligand in the CRC cell lines. ER $\beta$  has previously been shown to have ligand-independent functions in cell lines, which may be a consequence of overexpression or that the high level of growth factors activates ER $\beta$  through phosphorylation. Hence, we did not treat the cells with E2 to activate ER $\beta$ . Furthermore, the addition of TNF $\alpha$  to estrogen (E2) treatment in the MCF-7 breast cancer cells has been reported to alter the ER $\alpha$  enhancer landscape in an NF $\kappa$ B-dependent manner (275). In future studies, it would be interesting to investigate whether the ER $\beta$  cistrome is affected by TNF $\alpha$  treatment. Here, we defined how ER $\beta$  modulated p65 chromatin binding in colon cancer cells and provided the mechanistic basis for understanding the p65 cistrome and corresponding TNF $\alpha$ -induced response; further revealing a cross-talk between ER $\beta$  and the NF $\kappa$ B pathway (Figure 9).



**Figure 9.** Schematic summary model of ER $\beta$  genomic binding and cross-talk with potential transcriptional factors such as p65, AP-1 and TCF motifs and the regulated genes are involved in inflammatory and cellular processes which are important in CRC formation and development.

In conclusion, our data indicate that intestinal ER $\beta$  is protective against CA-CRC in a sex-dependent manner by attenuating the TNF $\alpha$ -NF $\kappa$ B pathway. Through genome-wide binding studies and transcriptomic studies, we demonstrate that ER $\beta$  regulates several tumor suppressors and oncogenes involved in CRC progression. We identified that ER $\beta$  impacts the p65 cistrome and, in particular, regulates circadian clock genes and modulates p65 chromatin binding to circadian clock genes. Estrogen through ER $\beta$  also protects against HFD-induced deregulation and impacts epithelial cell proliferation and macrophage infiltration in a sex-dependent manner.

## 5 CONCLUSIONS

The following are the major findings of this research:

- Loss of intestinal ER $\beta$  enhances CRC in a sex-dependent manner, increasing the incidence of tumors in males and tumor size in females. This suggests that ER $\beta$ , which is present in the intestinal epithelial cells, is protective against CRC.
- ER $\beta$  attenuates TNF $\alpha$ -induced epithelial cell damage and reduces inflammation, as well as impacts intestinal crypt cell proliferation.
- In colon cancer cells, ER $\beta$  binds and regulates several important tumor suppressors and oncogenes involved in the oncogenic Wnt/ $\beta$ -catenin pathway and NF $\kappa$ B signaling. Novel major motifs identified in ER $\beta$  chromatin binding sites include TCF and KLF motifs. This indicates potential genomic cross-talk between ER $\beta$  and TCF and KLF factors.
- Estrogen via ER $\beta$  protects against the HFD-induced pro-inflammatory condition. Selective activation of ER $\beta$  with DPN reduced macrophage infiltration and epithelial cell proliferation. Intestinal ER $\beta$  restored the HFD dysregulated clock genes expression *in vivo*.
- ER $\beta$  attenuates the TNF $\alpha$  transcriptome in CRC cell lines and modifies p65 chromatin binding in a cell line-dependent manner. It also modulates the expression of NF $\kappa$ B target genes. ER $\beta$  impacts p65 chromatin binding to core clock genes in CRC cell lines.
- The circadian clock is essential to maintain gut function, inflammation, and metabolism, and ER $\beta$  regulating this may be imperative for its protective activity. This indicates a key role for ER $\beta$  in the regulation of clock genes, providing a novel approach to study the impact of ER $\beta$  in colonic circadian rhythm.

## 6 FUTURE PERSPECTIVES

The studies conducted with our animal model ER $\beta$ KO<sup>Vil</sup> suggest that loss of intestinal ER $\beta$  aggravates CRC and enhances TNF $\alpha$  signaling. We recently also performed RNA-sequencing of WT and ER $\beta$ KO<sup>Vil</sup> colon to investigate the protective effects of ER $\beta$ . In the future, it would be interesting to perform both AOM-DSS and HFD studies in a recently developed novel transgenic mouse model Gt(ROSA)26Sor with ER $\beta$  overexpression (iER $\beta$ +). Endogenous ER $\beta$  expression is very low in the colon, and it would be fascinating to compare the results with the ER $\beta$  knock-in model. This would help advance our understanding of the localized impact of intestinal ER $\beta$  on colon microenvironment during colitis and HFD-induced obesity. Future studies should also focus on the effects of restoring ER $\beta$  expression in adenomas (pre-stage of CRC) using an inducible intestinal ER $\beta$  mouse model. This would add to our understanding of whether restoring ER $\beta$  in tumors could exert anti-tumorigenic effects. Further, investigating approaches to restore ER $\beta$  expression in the advanced stages of CRC, as has been reported in polyps by certain phytoestrogenic/lignan complexes. Several doxycycline-inducible mouse models available are not suitable for colon studies, as doxycycline impacts the gut microbiome by decreasing bacterial diversity, richness (346), and metabolism (347). Developing safe inducible intestinal ER $\beta$  mouse models could enable us to further explore ER $\beta$  activity in detail.

My cistrome data identified novel evidence of interaction between ER $\beta$  and TCF and KLF motifs. More insight into this cross-talk can be achieved from CO-IP or possibly IP-MS. Recent methodological developments such as RIME (rapid immunoprecipitation mass spectrometry of endogenous proteins) or quantitative multiplexed RIME methods further help to explore protein interactions in complex chromatin heterogeneity (348, 349). Simultaneously, exploring ER $\beta$  interaction with AP1 or nuclear receptors expressed in colon is also essential to target NR cross-talk. In the future, this would help to design ligands to improve or restrict transcription factor cross-talk. So far, due to low levels of ER $\beta$  expression in the colon, it is challenging to perform ChIP-Seq using endogenous expression of ER $\beta$ . The iER $\beta$ + mouse would also provide an opportunity to further corroborate the ChIP-Seq results *in vivo*.

In this study, we sequenced E2-activated ER $\beta$  ChIP samples. However, we also demonstrated (at a few selected sites) that the DNA binding of ER $\beta$  is ligand-independent in these cell lines. The reason for this is not known. In COS1 cells, it has been shown that EGF (epidermal growth factor) induces phosphorylation of serine residues (S104, S124) of ER $\beta$  and activates its interaction with steroidal co-receptor activator 1 (SRC-1), and this results in ligand-independent activity (350, 351). However, we cannot rule out the possibility that the E2 independence results from ER $\beta$  overexpression. In this regard, it is also interesting to perform ChIP-Seq without E2 ligand and compare both cistromes to determine if there is any difference present, in some or a proportion of binding sites, between ligand-activated ER $\beta$  and ligand-independent ER $\beta$  binding patterns.

In the HFD study, we have studied how ER $\beta$  impacts HFD-induced dysregulation of clock genes at only one time point. Further analyzing this at different Zeitgeber time points throughout the day will help to identify how ER $\beta$  affects circadian rhythmicity. We observed substantial sex differences in the colon transcriptome of mice. Another study from our group recently identified sexual dimorphism in humans in normal colon and colon cancer transcriptomes and presented sex-specific prognostic CRC biomarkers (352). Such sex differences may help to explain the difference in CRC incidence, tumor characteristics, and mortality between men and women. Altogether, it appears imperative that future studies of CRC and colon consider sex in their design.

My Ph.D. studies have made an interesting finding that p65 binding to circadian clock genes in colon cancer cells. More insights into this regulation could help to understand this novel finding. I have revealed that the ER $\beta$  impact on p65 activity is cell-type dependent. Also, our data shows an interesting feedback loop mechanism in which TNF $\alpha$  transactivates ER $\beta$ , which in turn inhibits TNF $\alpha$  signaling. Therefore, further research into TNF $\alpha$ -P65-ER $\beta$  cross-talk, *in vitro* and *in vivo*, is required.

## 7 ACKNOWLEDGEMENTS

I am grateful to everyone who directly or indirectly contributed to my thesis.

First, I would like to express my gratitude to my supervisor, **Cecilia Williams**, for allowing me to be a part of an excellent team working on exciting projects. Your valuable advice and constructive criticism guided my development as an independent scientist. You are always cheerful and smiling, from our first meeting to the last one we had yesterday. I always left our meetings with a sense of accomplishment and motivation. You helped me a lot to make this long Ph.D. journey incredible and knowledgeable.

**Amena Archer**, for your immense support in every project. Thank you for the constructive discussions and for including me in many collaborations from which I learned a lot. I remember the late nights we spent in the lab and the phone calls during this pandemic. You influenced me greatly and showed me the value of life skills. I would like to thank **Afshin Ahmadian**, my co-supervisor, and **Karin Dahlman-Wright**, previous co-supervisor, for your support and good advice.

On this note, I thank my fellow PhD-fighter, **Linnea Hases**, for being a huge support in the lab for valuable discussions, and I truly look up to the vision you have towards work. I treasure the times we spent together in Greece. You are the person who made me realize the impossible is possible, well that is making me learn how to swim. I want to thank **Mohammed Hakim Jaffer Ali** for being a big brother in the lab and helping me in the cell lab. **Madeleine Birgersson** for all the night meetings in the animal facility, good scientific discussions, and talks over lunch. Thanks for being my Swedish translator. **Dandan Song**, it was fun to work with you and your kind support when I need something in the lab. **Fahmi Mesmar** and **Ahmed Ibrahim**, for your support in the beginning. Thank you to all the master students who helped contribute to the work.

I want to thank my mentor **Anastasios Damdimopoulos**, for being a huge pillar of support in data analysis and teaching me a lot about ChIP-Seq. I also thank everyone in BEA (Bioinformatics and Expression Analysis) for ChIP-Sequencing. Thanks to **Christian Oretlin** and **Max Karlsson** for your assistance in R in the beginning and **Johannes Ristau** for help in the lab.

**Monica Ahlberg** and **Karin Gåse** at the KI administration, thank you for all your assistance with my long application forms.

Thank you to the members of my half-time examination board, **Eckardt Treuter**, **Monika Nistér**, and **Jochen Schwenk**, for valuable advice during my half-time.

Thanks to **Jason Carroll** for being my opponent and **Ann-Kristin Östlund Farrants**, **Eckardt Treuter**, and **Monika Nistér** for taking part in the examination board.

My Friends in Sweden, **Arun, Yasamin, Tejaswi, and Sowjanya**, thanks for the beautiful times we spent together. **Poonam and Monil**, Thanks for all the fun times and movie nights that kept me active. **Teja and Bhanu**, Thanks for all the beautiful moments we shared over the weekend lunch meetings. Special thanks to **Saritha and Sultana** for their nice company and kind support.

I have to express my gratitude to my husband, **Anand Varma**, for his understanding and support. Thank you for the many years of joy and inspiration you have given me. Heartfelt thanks to my wonderful daughter **Aashira** who keep me grounded with her charm and happiness.

Thanks to my parents **Nagaraju and Varalakshmi**, for their love and constant support throughout my life. Thanks, mom, who always given me the strength to chase my dreams. My dad always inspires me to keep working hard with utmost determination and passion towards what we are doing. Thank both **Vijaya Gaaru and Raju Gaaru** for your guidance and constant support in this long journey. I would also like to thank my sibling **Sai** for being a great, supportive brother. Thanks to my beloved grandparents **Krishnaveni and Subba Raju** for their love and encouragement in all my endeavors. **Pushpa**, thanks for always being there for me. **Jyothi, Anantha, and Geetha**, for your help and advice during my graduation time. My two close friends and brothers, **Narasimha Raju and Swamy**, will always be remembered.

A huge thanks to my family members, my aunts **Jyothi and Rajeshwari**, who traveled from India to Sweden to spend time with my family. Thanks to my wonderful group of cousins **Nimisha, Vinay, Manoj, Harika, Santhi, Padmaja, Sathvik, Charmi, Teju, and Chinnu** for all the happy moments and inspirations.

On this note, I take the privilege to thank two of my best buddies for fifteen years **Sowjanya and Abhi**, far away in person, yet so close by hearts. You are the people who stood by me through thick and thin. Thank you!



## 8 REFERENCES

1. Chapelle N, Martel M, Toes-Zoutendijk E, Barkun AN, Bardou M. Recent advances in clinical practice: colorectal cancer chemoprevention in the average-risk population. *Gut*. 2020;69(12):2244-55.
2. Schneider LS, Hernandez G, Zhao L, Franke AA, Chen Y-L, Pawluczyk S, et al. Safety and feasibility of estrogen receptor- $\beta$  targeted phytoSERM formulation for menopausal symptoms: phase 1b/2a randomized clinical trial. *Menopause*. 2019;26(8).
3. Tagliaferri MA, Tagliaferri MC, Creasman JM, Koltun WD. A Selective Estrogen Receptor Beta Agonist for the Treatment of Hot Flushes: Phase 2 Clinical Trial. *J Altern Complement Med*. 2016;22(9):722-8.
4. Kuipers EJ, Rösch T, Bretthauer M. Colorectal cancer screening—optimizing current strategies and new directions. *Nature Reviews Clinical Oncology*. 2013;10:130.
5. Kuipers EJ, Grady WM, Lieberman D, Seufferlein T, Sung JJ, Boelens PG, et al. Colorectal cancer. *Nature reviews Disease primers*. 2015;1:15065-.
6. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA: A Cancer Journal for Clinicians*. 2018;68(1):7-30.
7. Arnold M, Sierra MS, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global patterns and trends in colorectal cancer incidence and mortality. *Gut*. 2017;66(4):683-91.
8. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018;68(6):394-424.
9. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin*. 2019;69(1):7-34.
10. Fidler MM, Soerjomataram I, Bray F. A global view on cancer incidence and national levels of the human development index. *International journal of cancer*. 2016;139(11):2436-46.
11. Bray F. Transition in human development and the global cancer burden. *World cancer report 2014*. 2014:54-68.
12. Siegel RL, Miller KD, Goding Sauer A, Fedewa SA, Butterly LF, Anderson JC, et al. Colorectal cancer statistics, 2020. *CA Cancer J Clin*. 2020;70(3):145-64.
13. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: A Cancer Journal for Clinicians*. n/a(n/a).
14. Kasi PM, Shahjehan F, Cochuyt JJ, Li Z, Colibaseanu DT, Merchea A. Rising Proportion of Young Individuals With Rectal and Colon Cancer. *Clinical Colorectal Cancer*. 2019;18(1):e87-e95.
15. Siegel RL, Torre LA, Soerjomataram I, Hayes RB, Bray F, Weber TK, et al. Global patterns and trends in colorectal cancer incidence in young adults. *Gut*. 2019;68(12):2179.
16. Vuik FE, Nieuwenburg SA, Bardou M, Lansdorp-Vogelaar I, Dinis-Ribeiro M, Bento MJ, et al. Increasing incidence of colorectal cancer in young adults in Europe over the last 25 years. *Gut*. 2019;68(10):1820-6.

17. Thrumurthy SG, Thrumurthy SSD, Gilbert CE, Ross P, Haji A. Colorectal adenocarcinoma: risks, prevention and diagnosis. *BMJ*. 2016;354:i3590.
18. Galiatsatos P, Foulkes WD. Familial adenomatous polyposis. *Am J Gastroenterol*. 2006;101(2):385-98.
19. Yurgelun MB, Kulke MH, Fuchs CS, Allen BA, Uno H, Hornick JL, et al. Cancer Susceptibility Gene Mutations in Individuals With Colorectal Cancer. *J Clin Oncol*. 2017;35(10):1086-95.
20. Soderlund S, Granath F, Brostrom O, Karlen P, Lofberg R, Ekbom A, et al. Inflammatory bowel disease confers a lower risk of colorectal cancer to females than to males. *Gastroenterology*. 2010;138(5):1697-703.
21. Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA, Jr., Kinzler KW. Cancer genome landscapes. *Science (New York, NY)*. 2013;339(6127):1546-58.
22. Platz EA, Willett WC, Colditz GA, Rimm EB, Spiegelman D, Giovannucci E. Proportion of colon cancer risk that might be preventable in a cohort of middle-aged US men. *Cancer Causes Control*. 2000;11(7):579-88.
23. Erdrich J, Zhang X, Giovannucci E, Willett W. Proportion of colon cancer attributable to lifestyle in a cohort of US women. *Cancer Causes Control*. 2015;26(9):1271-9.
24. Kune GA, Bannerman S, Watson LF. Attributable risk for diet, alcohol, and family history in the Melbourne colorectal cancer study. *Nutrition and Cancer*. 1992;18(3):231-5.
25. Chan DS, Lau R, Aune D, Vieira R, Greenwood DC, Kampman E, et al. Red and processed meat and colorectal cancer incidence: meta-analysis of prospective studies. *PLoS One*. 2011;6(6):e20456.
26. Cotterchio M, Boucher BA, Manno M, Gallinger S, Okey A, Harper P. Dietary phytoestrogen intake is associated with reduced colorectal cancer risk. *The Journal of nutrition*. 2006;136(12):3046-53.
27. Fuchs MA, Yuan C, Sato K, Niedzwiecki D, Ye X, Saltz LB, et al. Predicted vitamin D status and colon cancer recurrence and mortality in CALGB 89803 (Alliance). *Annals of oncology : official journal of the European Society for Medical Oncology*. 2017;28(6):1359-67.
28. Keum N, Aune D, Greenwood DC, Ju W, Giovannucci EL. Calcium intake and colorectal cancer risk: dose-response meta-analysis of prospective observational studies. *International journal of cancer*. 2014;135(8):1940-8.
29. Parkin DM. 9. Cancers attributable to inadequate physical exercise in the UK in 2010. *British Journal Of Cancer*. 2011;105:S38.
30. Wolin KY, Yan Y, Colditz GA, Lee IM. Physical activity and colon cancer prevention: a meta-analysis. *British Journal Of Cancer*. 2009;100:611.
31. Amin MB, Greene FL, Edge SB, Compton CC, Gershenwald JE, Brookland RK, et al. The Eighth Edition AJCC Cancer Staging Manual: Continuing to build a bridge from a population-based to a more "personalized" approach to cancer staging. *CA Cancer J Clin*. 2017;67(2):93-9.
32. Trinh A, Ladrach C, Dawson HE, ten Hoorn S, Kuppen PJK, Reimers MS, et al. Tumour budding is associated with the mesenchymal colon cancer subtype and RAS/RAF mutations: a study of 1320 colorectal cancers with Consensus Molecular Subgroup (CMS) data. *British Journal of Cancer*. 2018;119(10):1244-51.

33. Guinney J, Dienstmann R, Wang X, de Reyniès A, Schlicker A, Sonesson C, et al. The consensus molecular subtypes of colorectal cancer. *Nature Medicine*. 2015;21(11):1350-6.
34. Navarro M, Nicolas A, Ferrandez A, Lanás A. Colorectal cancer population screening programs worldwide in 2016: An update. *World J Gastroenterol*. 2017;23(20):3632-42.
35. Blom J, Kilpeläinen S, Hultcrantz R, Törnberg S. Five-year experience of organized colorectal cancer screening in a Swedish population – increased compliance with age, female gender, and subsequent screening round. *Journal of Medical Screening*. 2014;21(3):144-50.
36. Ruder EH, Laiyemo AO, Graubard BI, Hollenbeck AR, Schatzkin A, Cross AJ. Non-steroidal anti-inflammatory drugs and colorectal cancer risk in a large, prospective cohort. *The American journal of gastroenterology*. 2011;106(7):1340-50.
37. Flossmann E, Rothwell PM. Effect of aspirin on long-term risk of colorectal cancer: consistent evidence from randomised and observational studies. *Lancet (London, England)*. 2007;369(9573):1603-13.
38. Arber N. Cyclooxygenase-2 Inhibitors in Colorectal Cancer Prevention: Point. *Cancer Epidemiology Biomarkers & Prevention*. 2008;17(8):1852-7.
39. Bresalier RS, Sandler RS, Quan H, Bolognese JA, Oxenius B, Horgan K, et al. Cardiovascular events associated with rofecoxib in a colorectal adenoma chemoprevention trial. *The New England journal of medicine*. 2005;352(11):1092-102.
40. Zheng SL, Roddick AJ. Association of Aspirin Use for Primary Prevention With Cardiovascular Events and Bleeding Events: A Systematic Review and Meta-analysis Association of Aspirin Use for Primary Prevention of CVD With Cardiovascular Events and Bleeding Association of Aspirin Use for Primary Prevention of CVD With Cardiovascular Events and Bleeding. *JAMA*. 2019;321(3):277-87.
41. Nolfo F, Rametta S, Marventano S, Grosso G, Mistretta A, Drago F, et al. Pharmacological and dietary prevention for colorectal cancer. *BMC Surg*. 2013;13 Suppl 2(Suppl 2):S16-S.
42. Routine aspirin or nonsteroidal anti-inflammatory drugs for the primary prevention of colorectal cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2007;146(5):361-4.
43. Xie Y-H, Chen Y-X, Fang J-Y. Comprehensive review of targeted therapy for colorectal cancer. *Signal Transduction and Targeted Therapy*. 2020;5(1):22.
44. Grady WM, Carethers JM. Genomic and epigenetic instability in colorectal cancer pathogenesis. *Gastroenterology*. 2008;135(4):1079-99.
45. Boland CR, Goel A. Microsatellite Instability in Colorectal Cancer. *Gastroenterology*. 2010;138(6):2073-87.e3.
46. Weisenberger DJ, Siegmund KD, Campan M, Young J, Long TI, Faasse MA, et al. CpG island methylator phenotype underlies sporadic microsatellite instability and is tightly associated with BRAF mutation in colorectal cancer. *Nature genetics*. 2006;38(7):787-93.
47. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 2011;144(5):646-74.
48. Basu S, Haase G, Ben-Ze'ev A. Wnt signaling in cancer stem cells and colon cancer metastasis. *F1000Res*. 2016;5:F1000 Faculty Rev-699.

49. Kemper K, Prasetyanti PR, De Lau W, Rodermond H, Clevers H, Medema JP. Monoclonal antibodies against Lgr5 identify human colorectal cancer stem cells. *Stem Cells*. 2012;30(11):2378-86.
50. Vermeulen L, De Sousa E Melo F, van der Heijden M, Cameron K, de Jong JH, Borovski T, et al. Wnt activity defines colon cancer stem cells and is regulated by the microenvironment. *Nature cell biology*. 2010;12(5):468-76.
51. Miyoshi Y, Nagase H, Ando H, Horii A, Ichii S, Nakatsuru S, et al. Somatic mutations of the APC gene in colorectal tumors: mutation cluster region in the APC gene. *Hum Mol Genet*. 1992;1(4):229-33.
52. Herbst A, Jurinovic V, Krebs S, Thieme SE, Blum H, Göke B, et al. Comprehensive analysis of  $\beta$ -catenin target genes in colorectal carcinoma cell lines with deregulated Wnt/ $\beta$ -catenin signaling. *BMC Genomics*. 2014;15(1):74.
53. Liu J, Xing Y, Hinds TR, Zheng J, Xu W. The third 20 amino acid repeat is the tightest binding site of APC for beta-catenin. *Journal of molecular biology*. 2006;360(1):133-44.
54. Sparks AB, Morin PJ, Vogelstein B, Kinzler KW. Mutational Analysis of the APC/ $\beta$ -Catenin/Tcf Pathway in Colorectal Cancer. *Cancer Research*. 1998;58(6):1130-4.
55. Malumbres M, Barbacid M. RAS oncogenes: the first 30 years. *Nature Reviews Cancer*. 2003;3:459.
56. Morikawa T, Kuchiba A, Liao X, Imamura Y, Yamauchi M, Qian ZR, et al. Tumor TP53 expression status, body mass index and prognosis in colorectal cancer. *International journal of cancer*. 2012;131(5):1169-78.
57. Colussi D, Brandi G, Bazzoli F, Ricciardiello L. Molecular pathways involved in colorectal cancer: implications for disease behavior and prevention. *International journal of molecular sciences*. 2013;14(8):16365-85.
58. Massagué J. TGF $\beta$  in Cancer. *Cell*. 2008;134(2):215-30.
59. Fleming NI, Jorissen RN, Mouradov D, Christie M, Sakthianandeswaren A, Palmieri M, et al. SMAD2, SMAD3 and SMAD4 mutations in colorectal cancer. *Cancer Res*. 2013;73(2):725-35.
60. Xia L, Tan S, Zhou Y, Lin J, Wang H, Oyang L, et al. Role of the NF $\kappa$ B-signaling pathway in cancer. *OncoTargets and therapy*. 2018;11:2063-73.
61. Karin M, Cao Y, Greten FR, Li Z-W. NF- $\kappa$ B in cancer: from innocent bystander to major culprit. *Nature Reviews Cancer*. 2002;2:301.
62. Haefner B. NF- $\kappa$ B: arresting a major culprit in cancer. *Drug Discovery Today*. 2002;7(12):653-63.
63. Rayet B, G  linas C. Aberrant rel/nfkb genes and activity in human cancer. *Oncogene*. 1999;18:6938.
64. Serasanambati M, Chilakapati SR. Function of Nuclear Factor Kappa B (NF- $\kappa$ B) in human diseases- A Review2016. 368 p.
65. Singh N, Baby D, Rajguru JP, Patil PB, Thakkannavar SS, Pujari VB. Inflammation and cancer. *Ann Afr Med*. 2019;18(3):121-6.

66. O'Connor PM, Lapointe TK, Beck PL, Buret AG. Mechanisms by which inflammation may increase intestinal cancer risk in inflammatory bowel disease. *Inflammatory Bowel Diseases*. 2010;16(8):1411-20.
67. Souici AC, Mirkovitch J, Hausel P, Keefer LK, Felley-Bosco E. Transition mutation in codon 248 of the p53 tumor suppressor gene induced by reactive oxygen species and a nitric oxide-releasing compound. *Carcinogenesis*. 2000;21(2):281-7.
68. Ambs S, Bennett WP, Merriam WG, Ogunfusika MO, Oser SM, Harrington AM, et al. Relationship between p53 mutations and inducible nitric oxide synthase expression in human colorectal cancer. *J Natl Cancer Inst*. 1999;91(1):86-8.
69. Ricciardi M, Zanotto M, Malpeli G, Bassi G, Perbellini O, Chilosi M, et al. Epithelial-to-mesenchymal transition (EMT) induced by inflammatory priming elicits mesenchymal stromal cell-like immune-modulatory properties in cancer cells. *British Journal of Cancer*. 2015;112(6):1067-75.
70. Bian XW, Chen JH, Jiang XF, Bai JS, Wang QL, Zhang X. Angiogenesis as an immunopharmacologic target in inflammation and cancer. *International immunopharmacology*. 2004;4(12):1537-47.
71. Sandler RS, Halabi S, Baron JA, Budinger S, Paskett E, Keresztes R, et al. A Randomized Trial of Aspirin to Prevent Colorectal Adenomas in Patients with Previous Colorectal Cancer. *New England Journal of Medicine*. 2003;348(10):883-90.
72. Din FVN, Theodoratou E, Farrington SM, Tenesa A, Barnetson RA, Cetnarskyj R, et al. Effect of aspirin and NSAIDs on risk and survival from colorectal cancer. *Gut*. 2010;59(12):1670-9.
73. Nishihara R, Lochhead P, Kuchiba A, Jung S, Yamauchi M, Liao X, et al. Aspirin use and risk of colorectal cancer according to BRAF mutation status. *Jama*. 2013;309(24):2563-71.
74. Rosen MJ, Dhawan A, Saeed SA. Inflammatory Bowel Disease in Children and Adolescents. *JAMA Pediatr*. 2015;169(11):1053-60.
75. Gramlich T, Petras RE. Pathology of inflammatory bowel disease. *Seminars in Pediatric Surgery*. 2007;16(3):154-63.
76. Choi PM, Zelig MP. Similarity of colorectal cancer in Crohn's disease and ulcerative colitis: implications for carcinogenesis and prevention. *Gut*. 1994;35(7):950-4.
77. Kim BJ, Yang SK, Kim JS, Jeon YT, Choi H, Han DS, et al. Trends of ulcerative colitis-associated colorectal cancer in Korea: A KASID study. *J Gastroenterol Hepatol*. 2009;24(4):667-71.
78. Kameyama H, Nagahashi M, Shimada Y, Tajima Y, Ichikawa H, Nakano M, et al. Genomic characterization of colitis-associated colorectal cancer. *World J Surg Oncol*. 2018;16(1):121-.
79. Ning Y, Wang L, Giovannucci EL. A quantitative analysis of body mass index and colorectal cancer: findings from 56 observational studies. *Obesity Reviews*. 2010;11(1):19-30.
80. Guilherme A, Virbasius JV, Puri V, Czech MP. Adipocyte dysfunctions linking obesity to insulin resistance and type 2 diabetes. *Nat Rev Mol Cell Biol*. 2008;9(5):367-77.
81. Poritz LS, Garver KI, Green C, Fitzpatrick L, Ruggiero F, Koltun WA. Loss of the tight junction protein ZO-1 in dextran sulfate sodium induced colitis. *J Surg Res*. 2007;140(1):12-9.

82. Cani PD, Bibiloni R, Knauf C, Waget A, Neyrinck AM, Delzenne NM, et al. Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. *Diabetes*. 2008;57(6):1470-81.
83. Kawano Y, Nakae J, Watanabe N, Kikuchi T, Tateya S, Tamori Y, et al. Colonic Pro-inflammatory Macrophages Cause Insulin Resistance in an Intestinal Ccl2/Ccr2-Dependent Manner. *Cell Metabolism*. 2016;24(2):295-310.
84. Fu T, Coulter S, Yoshihara E, Oh TG, Fang S, Cayabyab F, et al. FXR Regulates Intestinal Cancer Stem Cell Proliferation. *Cell*. 2019;176(5):1098-112.e18.
85. Zarrinpar A, Chaix A, Panda S. Daily Eating Patterns and Their Impact on Health and Disease. *Trends in Endocrinology & Metabolism*. 2016;27(2):69-83.
86. Cunningham PS, Ahern SA, Smith LC, da Silva Santos CS, Wager TT, Bechtold DA. Targeting of the circadian clock via CK1 $\delta/\epsilon$  to improve glucose homeostasis in obesity. *Scientific Reports*. 2016;6(1):29983.
87. Kohsaka A, Laposky AD, Ramsey KM, Estrada C, Joshu C, Kobayashi Y, et al. High-Fat Diet Disrupts Behavioral and Molecular Circadian Rhythms in Mice. *Cell Metabolism*. 2007;6(5):414-21.
88. Sun L, Wang Y, Song Y, Cheng X-R, Xia S, Rahman MRT, et al. Resveratrol restores the circadian rhythmic disorder of lipid metabolism induced by high-fat diet in mice. *Biochemical and Biophysical Research Communications*. 2015;458(1):86-91.
89. Voigt RM, Forsyth CB, Keshavarzian A. Circadian rhythms: a regulator of gastrointestinal health and dysfunction. *Expert Rev Gastroenterol Hepatol*. 2019;13(5):411-24.
90. Sun M, Feng W, Wang F, Li P, Li Z, Li M, et al. Meta-analysis on shift work and risks of specific obesity types. *Obes Rev*. 2018;19(1):28-40.
91. Wang X, Ji A, Zhu Y, Liang Z, Wu J, Li S, et al. A meta-analysis including dose-response relationship between night shift work and the risk of colorectal cancer. *Oncotarget*. 2015;6(28).
92. Kim S-E, Paik HY, Yoon H, Lee JE, Kim N, Sung M-K. Sex- and gender-specific disparities in colorectal cancer risk. *World J Gastroenterol*. 2015;21(17):5167-75.
93. Wang G, Wang J-J, Yin P-H, Xu K, Wang Y-Z, Shi F, et al. New strategies for targeting glucose metabolism-mediated acidosis for colorectal cancer therapy. *Journal of Cellular Physiology*. 2019;234(1):348-68.
94. Brown DG, Rao S, Weir TL, O'Malia J, Bazan M, Brown RJ, et al. Metabolomics and metabolic pathway networks from human colorectal cancers, adjacent mucosa, and stool. *Cancer Metab*. 2016;4:11.
95. Hall E, Volkov P, Dayeh T, Esguerra JLS, Salö S, Eliasson L, et al. Sex differences in the genome-wide DNA methylation pattern and impact on gene expression, microRNA levels and insulin secretion in human pancreatic islets. *Genome biology*. 2014;15(12):522.
96. Jia M, Dahlman-Wright K, Gustafsson J-Å. Estrogen receptor alpha and beta in health and disease. *Best practice & research Clinical endocrinology & metabolism*. 2015;29(4):557-68.
97. Chen C, Gong X, Yang X, Shang X, Du Q, Liao Q, et al. The roles of estrogen and estrogen receptors in gastrointestinal disease. *Oncol Lett*. 2019;18(6):5673-80.
98. Kim YS, Kim N, Kim GH. Sex and Gender Differences in Gastroesophageal Reflux Disease. *J Neurogastroenterol Motil*. 2016;22(4):575-88.

99. Meleine M, Matricon J. Gender-related differences in irritable bowel syndrome: potential mechanisms of sex hormones. *World J Gastroenterol*. 2014;20(22):6725-43.
100. Cui J, Shen Y, Li R. Estrogen synthesis and signaling pathways during aging: from periphery to brain. *Trends in Molecular Medicine*. 2013;19(3):197-209.
101. Hatcher KM, Royston SE, Mahoney MM. Modulation of circadian rhythms through estrogen receptor signaling. *Eur J Neurosci*. 2020;51(1):217-28.
102. Bondesson M, Hao R, Lin C-Y, Williams C, Gustafsson J-Å. Estrogen receptor signaling during vertebrate development. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms*. 2015;1849(2):142-51.
103. Fuentes N, Silveyra P. Estrogen receptor signaling mechanisms. *Adv Protein Chem Struct Biol*. 2019;116:135-70.
104. Ruiz-Cortés ZT. Gonadal Sex Steroids: Production, Action and Interactions in Mammals 2012.
105. Simpson ER. Sources of estrogen and their importance. *J Steroid Biochem Mol Biol*. 2003;86(3-5):225-30.
106. Barakat R, Oakley O, Kim H, Jin J, Ko CJ. Extra-gonadal sites of estrogen biosynthesis and function. *BMB Rep*. 2016;49(9):488-96.
107. Bulun SE, Zeitoun K, Sasano H, Simpson ER. Aromatase in aging women. *Semin Reprod Endocrinol*. 1999;17(4):349-58.
108. Holinka C, Diczfalussy E, Coelingh Bennink H. Estetrol: A unique steroid in human pregnancy. *The Journal of steroid biochemistry and molecular biology*. 2008;110:138-43.
109. Ali ES, Mangold C, Peiris AN. Estriol: emerging clinical benefits. *Menopause*. 2017;24(9):1081-5.
110. Watson CS, Jeng YJ, Kochukov MY. Nongenomic actions of estradiol compared with estrone and estriol in pituitary tumor cell signaling and proliferation. *Faseb j*. 2008;22(9):3328-36.
111. Kobayashi H, Yoshida S, Shirasawa N, Maeda K, Naito A. Expression and localization of aromatase in human gastric mucosa : Immunohistochemical study using biopsy materials. *Histochem Cell Biol*. 2019;151(1):21-8.
112. Kobayashi H. Estrogen synthesis in gastric parietal cells and secretion into portal vein. *Anatomical Science International*. 2020;95(1):22-30.
113. Dhiman VK, Bolt MJ, White KP. Nuclear receptors in cancer - uncovering new and evolving roles through genomic analysis. *Nature reviews Genetics*. 2018;19(3):160-74.
114. Rastinejad F, Huang P, Chandra V, Khorasanizadeh S. Understanding nuclear receptor form and function using structural biology. *Journal of molecular endocrinology*. 2013;51(3):T1-T21.
115. Mangelsdorf DJ, Thummel C, Beato M, Herrlich P, Schütz G, Umesono K, et al. The nuclear receptor superfamily: The second decade. *Cell*. 1995;83(6):835-9.
116. Fujita K, Nonomura N. Role of Androgen Receptor in Prostate Cancer: A Review. *World J Mens Health*. 2019;37(3):288-95.
117. Carroll JS. Mechanisms of oestrogen receptor (ER) gene regulation in breast cancer. *Eur J Endocrinol*. 2016;175(1):R41-R9.

118. Gertz J, Savic D, Varley KE, Partridge EC, Safi A, Jain P, et al. Distinct properties of cell-type-specific and shared transcription factor binding sites. *Molecular cell*. 2013;52(1):25-36.
119. Lo Sasso G, Bovenga F, Murzilli S, Salvatore L, Di Tullio G, Martelli N, et al. Liver X receptors inhibit proliferation of human colorectal cancer cells and growth of intestinal tumors in mice. *Gastroenterology*. 2013;144(7):1497-507, 507.e1-13.
120. Degirolamo C, Modica S, Palasciano G, Moschetta A. Bile acids and colon cancer: Solving the puzzle with nuclear receptors. *Trends in Molecular Medicine*. 2011;17(10):564-72.
121. Modica S, Gofflot F, Murzilli S, D'Orazio A, Salvatore L, Pellegrini F, et al. The intestinal nuclear receptor signature with epithelial localization patterns and expression modulation in tumors. *Gastroenterology*. 2010;138(2):636-48, 48.e1-12.
122. Jensen EV, Suzuki T, Kawashima T, Stumpf WE, Jungblut PW, DeSombre ER. A two-step mechanism for the interaction of estradiol with rat uterus. *Proceedings of the National Academy of Sciences of the United States of America*. 1968;59(2):632-8.
123. Enmark E, Peltö-Huikko M, Grandien K, Lagercrantz S, Lagercrantz J, Fried G, et al. Human estrogen receptor beta-gene structure, chromosomal localization, and expression pattern. *J Clin Endocrinol Metab*. 1997;82(12):4258-65.
124. Kuiper GG, Enmark E, Peltö-Huikko M, Nilsson S, Gustafsson JA. Cloning of a novel receptor expressed in rat prostate and ovary. *Proceedings of the National Academy of Sciences of the United States of America*. 1996;93(12):5925-30.
125. Koehler KF, Helguero LA, Haldosén L-A, Warner M, Gustafsson J-Ak. Reflections on the Discovery and Significance of Estrogen Receptor  $\beta$ . *Endocrine Reviews*. 2005;26(3):465-78.
126. Lewandowski S, Kalita K, Kaczmarek L. Estrogen receptor  $\beta$ : Potential functional significance of a variety of mRNA isoforms. *FEBS Lett*. 2002;524(1):1-5.
127. Porter BA, Ortiz MA, Bratslavsky G, Kotula L. Structure and Function of the Nuclear Receptor Superfamily and Current Targeted Therapies of Prostate Cancer. *Cancers*. 2019;11(12):1852.
128. Lannigan DA. Estrogen receptor phosphorylation. *Steroids*. 2003;68(1):1-9.
129. Rochette-Egly C. Nuclear receptors: integration of multiple signalling pathways through phosphorylation. *Cell Signal*. 2003;15(4):355-66.
130. Schwabe JW, Chapman L, Finch JT, Rhodes D. The crystal structure of the estrogen receptor DNA-binding domain bound to DNA: how receptors discriminate between their response elements. *Cell*. 1993;75(3):567-78.
131. Heldring N, Pike A, Andersson S, Matthews J, Cheng G, Hartman J, et al. Estrogen Receptors: How Do They Signal and What Are Their Targets. *Physiological Reviews*. 2007;87(3):905-31.
132. Tsai M-J, O'Malley BW. MOLECULAR MECHANISMS OF ACTION OF STEROID/THYROID RECEPTOR SUPERFAMILY MEMBERS. *Annual Review of Biochemistry*. 1994;63(1):451-86.
133. Nilsson S, Gustafsson J-Å. Estrogen Receptors: Therapies Targeted to Receptor Subtypes. *Clinical Pharmacology & Therapeutics*. 2011;89(1):44-55.



134. Kumar R, Zakharov MN, Khan SH, Miki R, Jang H, Toraldo G, et al. The dynamic structure of the estrogen receptor. *Journal of amino acids*. 2011;2011:812540.
135. Koide A, Zhao C, Naganuma M, Abrams J, Deighton-Collins S, Skafar DF, et al. Identification of regions within the F domain of the human estrogen receptor alpha that are important for modulating transactivation and protein-protein interactions. *Molecular endocrinology* (Baltimore, Md). 2007;21(4):829-42.
136. Paterni I, Granchi C, Katzenellenbogen JA, Minutolo F. Estrogen receptors alpha (ERalpha) and beta (ERbeta): subtype-selective ligands and clinical potential. *Steroids*. 2014;90:13-29.
137. Leung Y-K, Mak P, Hassan S, Ho S-M. Estrogen receptor (ER)- $\beta$  isoforms: A key to understanding ER- $\beta$  signaling. *Proceedings of the National Academy of Sciences*. 2006;103(35):13162-7.
138. Ogawa S, Inoue S, Watanabe T, Orimo A, Hosoi T, Ouchi Y, et al. Molecular cloning and characterization of human estrogen receptor betacx: a potential inhibitor of estrogen action in human. *Nucleic acids research*. 1998;26(15):3505-12.
139. Kim CK, Torcaso A, Asimes A, Chung WCJ, Pak TR. Structural and functional characteristics of oestrogen receptor  $\beta$  splice variants: Implications for the ageing brain. *J Neuroendocrinol*. 2018;30(2):10.1111/jne.12488.
140. Ciucci A, Ferrandina G, Mascilini F, Filippetti F, Scambia G, Zannoni GF, et al. Estrogen receptor  $\beta$ : Potential target for therapy in adult granulosa cell tumors? *Gynecol Oncol*. 2018;150(1):158-65.
141. Lewandowski S, Kalita K, Kaczmarek L. Estrogen receptor beta. Potential functional significance of a variety of mRNA isoforms. *FEBS Lett*. 2002;524(1-3):1-5.
142. Maruyama K, Endoh H, Sasaki-Iwaoka H, Kanou H, Shimaya E, Hashimoto S, et al. A novel isoform of rat estrogen receptor beta with 18 amino acid insertion in the ligand binding domain as a putative dominant negative regulator of estrogen action. *Biochem Biophys Res Commun*. 1998;246(1):142-7.
143. Zhao C, Toresson G, Xu L, Koehler KF, Gustafsson JA, Dahlman-Wright K. Mouse estrogen receptor beta isoforms exhibit differences in ligand selectivity and coactivator recruitment. *Biochemistry*. 2005;44(22):7936-44.
144. Donoghue LJ, Neufeld TI, Li Y, Arai Y, Coons LA, Korach KS. Differential Activation of a Mouse Estrogen Receptor  $\beta$  Isoform (mER $\beta$ 2) with Endocrine-Disrupting Chemicals (EDCs). *Environ Health Perspect*. 2017;125(4):634-42.
145. Kumar A, Bean LA, Rani A, Jackson T, Foster TC. Contribution of estrogen receptor subtypes, ER $\alpha$ , ER $\beta$ , and GPER1 in rapid estradiol-mediated enhancement of hippocampal synaptic transmission in mice. *Hippocampus*. 2015;25(12):1556-66.
146. Romano SN, Gorelick DA. Crosstalk between nuclear and G protein-coupled estrogen receptors. *Gen Comp Endocrinol*. 2018;261:190-7.
147. Barton M, Filardo EJ, Lolait SJ, Thomas P, Maggiolini M, Prossnitz ER. Twenty years of the G protein-coupled estrogen receptor GPER: Historical and personal perspectives. *The Journal of Steroid Biochemistry and Molecular Biology*. 2018;176:4-15.
148. Rosenbaum DM, Rasmussen SG, Kobilka BK. The structure and function of G-protein-coupled receptors. *Nature*. 2009;459(7245):356-63.

149. Revankar CM, Cimino DF, Sklar LA, Arterburn JB, Prossnitz ER. A Transmembrane Intracellular Estrogen Receptor Mediates Rapid Cell Signaling. *Science* (New York, NY). 2005;307(5715):1625-30.
150. Thomas P, Pang Y, Filardo EJ, Dong J. Identity of an Estrogen Membrane Receptor Coupled to a G Protein in Human Breast Cancer Cells. *Endocrinology*. 2005;146(2):624-32.
151. Filardo EJ, Quinn JA, Frackelton AR, Jr., Bland KI. Estrogen Action Via the G Protein-Coupled Receptor, GPR30: Stimulation of Adenylyl Cyclase and cAMP-Mediated Attenuation of the Epidermal Growth Factor Receptor-to-MAPK Signaling Axis. *Molecular Endocrinology*. 2002;16(1):70-84.
152. Prossnitz ER, Barton M. Estrogen biology: New insights into GPER function and clinical opportunities. *Molecular and cellular endocrinology*. 2014;389(1):71-83.
153. Morito K, Aomori T, Hirose T, Kinjo J, Hasegawa J, Ogawa S, et al. Interaction of phytoestrogens with estrogen receptors alpha and beta (II). *Biol Pharm Bull*. 2002;25(1):48-52.
154. Golden RJ, Noller KL, Titus-Ernstoff L, Kaufman RH, Mittendorf R, Stillman R, et al. Environmental endocrine modulators and human health: an assessment of the biological evidence. *Crit Rev Toxicol*. 1998;28(2):109-227.
155. Aravindakshan J, Gregory M, Marcogliese DJ, Fournier M, Cyr DG. Consumption of xenoestrogen-contaminated fish during lactation alters adult male reproductive function. *Toxicol Sci*. 2004;81(1):179-89.
156. Darbre PD. Metalloestrogens: an emerging class of inorganic xenoestrogens with potential to add to the oestrogenic burden of the human breast. *J Appl Toxicol*. 2006;26(3):191-7.
157. Manas ES, Xu ZB, Unwalla RJ, Somers WS. Understanding the selectivity of genistein for human estrogen receptor-beta using X-ray crystallography and computational methods. *Structure*. 2004;12(12):2197-207.
158. Mersereau JE, Levy N, Staub RE, Baggett S, Zogovic T, Chow S, et al. Liquiritigenin is a plant-derived highly selective estrogen receptor beta agonist. *Molecular and cellular endocrinology*. 2008;283(1-2):49-57.
159. Kuiper GGJM, Carlsson B, Grandien K, Enmark E, Häggblad J, Nilsson S, et al. Comparison of the Ligand Binding Specificity and Transcript Tissue Distribution of Estrogen Receptors  $\alpha$  and  $\beta$ . *Endocrinology*. 1997;138(3):863-70.
160. Brzozowski AM, Pike ACW, Dauter Z, Hubbard RE, Bonn T, Engström O, et al. Molecular basis of agonism and antagonism in the oestrogen receptor. *Nature*. 1997;389(6652):753-8.
161. Puranik NV, Srivastava P, Bhatt G, John Mary DJS, Limaye AM, Sivaraman J. Determination and analysis of agonist and antagonist potential of naturally occurring flavonoids for estrogen receptor (ER $\alpha$ ) by various parameters and molecular modelling approach. *Scientific Reports*. 2019;9(1):7450.
162. Paterni I, Bertini S, Granchi C, Macchia M, Minutolo F. Estrogen receptor ligands: a patent review update. *Expert Opinion on Therapeutic Patents*. 2013;23(10):1247-71.
163. Farooq A. Structural and Functional Diversity of Estrogen Receptor Ligands. *Curr Top Med Chem*. 2015;15(14):1372-84.

164. Potkul RK, Unger JM, Livingston RB, Crew KD, Wilczynski SP, Salomon CG, et al. Randomized trial of medroxyprogesterone acetate for the prevention of endometrial pathology from adjuvant tamoxifen for breast cancer: SWOG S9630. *NPJ Breast Cancer*. 2016;2:16024.
165. Killackey MA, Hakes TB, Pierce VK. Endometrial adenocarcinoma in breast cancer patients receiving antiestrogens. *Cancer Treat Rep*. 1985;69(2):237-8.
166. Stauffer SR, Coletta CJ, Tedesco R, Nishiguchi G, Carlson K, Sun J, et al. Pyrazole ligands: structure-affinity/activity relationships and estrogen receptor- $\alpha$ -selective agonists. *J Med Chem*. 2000;43(26):4934-47.
167. Meyers MJ, Sun J, Carlson KE, Marriner GA, Katzenellenbogen BS, Katzenellenbogen JA. Estrogen receptor- $\beta$  potency-selective ligands: structure-activity relationship studies of diarylpropionitriles and their acetylene and polar analogues. *J Med Chem*. 2001;44(24):4230-51.
168. Chadwick CC, Chippari S, Matelan E, Borges-Marcucci L, Eckert AM, Keith JC, Jr., et al. Identification of pathway-selective estrogen receptor ligands that inhibit NF- $\kappa$ B transcriptional activity. *Proceedings of the National Academy of Sciences of the United States of America*. 2005;102(7):2543-8.
169. Keith JC, Albert LM, Leathurby Y, Follettie M, Wang L, Borges-Marcucci L, et al. The utility of pathway selective estrogen receptor ligands that inhibit nuclear factor- $\kappa$ B transcriptional activity in models of rheumatoid arthritis. *Arthritis Research & Therapy*. 2005;7(3):R427.
170. Handa RJ, Weiser MJ, Zuloaga DG. A role for the androgen metabolite, 5 $\alpha$ -androstane-3 $\beta$ ,17 $\beta$ -diol, in modulating oestrogen receptor  $\beta$ -mediated regulation of hormonal stress reactivity. *J Neuroendocrinol*. 2009;21(4):351-8.
171. Hiroi R, Lacagnina AF, Hinds LR, Carbone DG, Uht RM, Handa RJ. The androgen metabolite, 5 $\alpha$ -androstane-3 $\beta$ ,17 $\beta$ -diol (3 $\beta$ -diol), activates the oxytocin promoter through an estrogen receptor- $\beta$  pathway. *Endocrinology*. 2013;154(5):1802-12.
172. Ho M-F, Correia C, Ingle JN, Kaddurah-Daouk R, Wang L, Kaufmann SH, et al. Ketamine and ketamine metabolites as novel estrogen receptor ligands: Induction of cytochrome P450 and AMPA glutamate receptor gene expression. *Biochemical Pharmacology*. 2018;152:279-92.
173. Klinge CM. Estrogen receptor interaction with estrogen response elements. *Nucleic acids research*. 2001;29(14):2905-19.
174. Bahadur U, Ganjam GK, Vasudevan N, Kondaiah P. Estrogen regulation of chicken riboflavin carrier protein gene is mediated by ERE half sites without direct binding of estrogen receptor. *Molecular and cellular endocrinology*. 2005;231(1):1-11.
175. Yaşar P, Ayaz G, User SD, Güpür G, Muyan M. Molecular mechanism of estrogen-estrogen receptor signaling. *Reprod Med Biol*. 2016;16(1):4-20.
176. Klinge CM. Estrogen receptor interaction with estrogen response elements. *Nucleic acids research*. 2001;29(14):2905-19.
177. Carleton JB, Ginley-Hidinger M, Berrett KC, Layer RM, Quinlan AR, Gertz J. Regulatory sharing between estrogen receptor  $\alpha$  bound enhancers. *Nucleic acids research*. 2020;48(12):6597-610.
178. Yi P, Driscoll MD, Huang J, Bhagat S, Hilf R, Bambara RA, et al. The effects of estrogen-responsive element- and ligand-induced structural changes on the recruitment of cofactors and

transcriptional responses by ER alpha and ER beta. *Molecular endocrinology* (Baltimore, Md). 2002;16(4):674-93.

179. Hyder SM, Chiappetta C, Stancel GM. Interaction of human estrogen receptors  $\alpha$  and  $\beta$  with the same naturally occurring estrogen response elements. *Biochemical Pharmacology*. 1999;57(6):597-601.

180. Mason CE, Shu FJ, Wang C, Session RM, Kallen RG, Sidell N, et al. Location analysis for the estrogen receptor-alpha reveals binding to diverse ERE sequences and widespread binding within repetitive DNA elements. *Nucleic acids research*. 2010;38(7):2355-68.

181. Lin CY, Vega VB, Thomsen JS, Zhang T, Kong SL, Xie M, et al. Whole-genome cartography of estrogen receptor alpha binding sites. *PLoS genetics*. 2007;3(6):e87.

182. Carroll JS, Liu XS, Brodsky AS, Li W, Meyer CA, Szary AJ, et al. Chromosome-wide mapping of estrogen receptor binding reveals long-range regulation requiring the forkhead protein FoxA1. *Cell*. 2005;122(1):33-43.

183. McDevitt MA, Glidewell-Kenney C, Jimenez MA, Ahearn PC, Weiss J, Jameson JL, et al. New insights into the classical and non-classical actions of estrogen: evidence from estrogen receptor knock-out and knock-in mice. *Molecular and cellular endocrinology*. 2008;290(1-2):24-30.

184. Safe S, Kim K. Non-classical genomic estrogen receptor (ER)/specificity protein and ER/activating protein-1 signaling pathways. *Journal of molecular endocrinology*. 2008;41(5):263-75.

185. Heldring N, Isaacs GD, Diehl AG, Sun M, Cheung E, Ranish JA, et al. Multiple sequence-specific DNA-binding proteins mediate estrogen receptor signaling through a tethering pathway. *Molecular endocrinology* (Baltimore, Md). 2011;25(4):564-74.

186. Jakacka M, Ito M, Martinson F, Ishikawa T, Lee EJ, Jameson JL. An estrogen receptor (ER)alpha deoxyribonucleic acid-binding domain knock-in mutation provides evidence for nonclassical ER pathway signaling in vivo. *Molecular endocrinology* (Baltimore, Md). 2002;16(10):2188-201.

187. Jakacka M, Ito M, Weiss J, Chien PY, Gehm BD, Jameson JL. Estrogen receptor binding to DNA is not required for its activity through the nonclassical AP1 pathway. *The Journal of biological chemistry*. 2001;276(17):13615-21.

188. Saville B, Hoivik D, Safe S, Porter W. Functional Synergy between the Transcription Factor Sp1 and the Estrogen Receptor. *Molecular Endocrinology*. 1997;11(11):1569-80.

189. Frasor J, Weaver A, Pradhan M, Dai Y, Miller LD, Lin C-Y, et al. Positive Cross-Talk between Estrogen Receptor and NF- $\kappa$ B in Breast Cancer. *Cancer Research*. 2009;69(23):8918-25.

190. Stender JD, Kim K, Charn TH, Komm B, Chang KCN, Kraus WL, et al. Genome-Wide Analysis of Estrogen Receptor  $\alpha$  DNA Binding and Tethering Mechanisms Identifies Runx1 as a Novel Tethering Factor in Receptor-Mediated Transcriptional Activation. *Molecular and Cellular Biology*. 2010;30(16):3943-55.

191. Wang X, Belguise K, Kersual N, Kirsch KH, Mineva ND, Galtier F, et al. Oestrogen signalling inhibits invasive phenotype by repressing RelB and its target BCL2. *Nature cell biology*. 2007;9(4):470-8.

192. Belguise K, Sonenshein GE. PKC $\theta$  promotes c-Rel-driven mammary tumorigenesis in mice and humans by repressing estrogen receptor  $\alpha$  synthesis. *The Journal of clinical investigation*. 2007;117(12):4009-21.
193. Frasor J, Weaver A, Pradhan M, Dai Y, Miller LD, Lin C-Y, et al. Positive cross-talk between estrogen receptor and NF-kappaB in breast cancer. *Cancer research*. 2009;69(23):8918-25.
194. Pradhan M, Baumgarten SC, Bembinster LA, Frasor J. CBP mediates NF- $\kappa$ B-dependent histone acetylation and estrogen receptor recruitment to an estrogen response element in the BIRC3 promoter. *Molecular and cellular biology*. 2012;32(2):569-75.
195. Zhao C, Gao H, Liu Y, Papoutsis Z, Jaffrey S, Gustafsson JA, et al. Genome-wide mapping of estrogen receptor-beta-binding regions reveals extensive cross-talk with transcription factor activator protein-1. *Cancer Res*. 2010;70(12):5174-83.
196. Paech K, Webb P, Kuiper GG, Nilsson S, Gustafsson J, Kushner PJ, et al. Differential ligand activation of estrogen receptors ERalpha and ERbeta at AP1 sites. *Science (New York, NY)*. 1997;277(5331):1508-10.
197. Björnström L, Sjöberg M. Mechanisms of Estrogen Receptor Signaling: Convergence of Genomic and Nongenomic Actions on Target Genes. *Molecular Endocrinology*. 2005;19(4):833-42.
198. Vanacker JM, Pettersson K, Gustafsson JA, Laudet V. Transcriptional targets shared by estrogen receptor-related receptors (ERRs) and estrogen receptor (ER)  $\alpha$ , but not by ERbeta. *Embo j*. 1999;18(15):4270-9.
199. Kim K, Thu N, Saville B, Safe S. Domains of estrogen receptor  $\alpha$  (ERalpha) required for ERalpha/Sp1-mediated activation of GC-rich promoters by estrogens and antiestrogens in breast cancer cells. *Molecular endocrinology (Baltimore, Md)*. 2003;17(5):804-17.
200. Anbalagan M, Rowan BG. Estrogen receptor  $\alpha$  phosphorylation and its functional impact in human breast cancer. *Molecular and cellular endocrinology*. 2015;418 Pt 3:264-72.
201. Siersbæk R, Kumar S, Carroll JS. Signaling pathways and steroid receptors modulating estrogen receptor  $\alpha$  function in breast cancer. *Genes Dev*. 2018;32(17-18):1141-54.
202. Simoncini T, Mannella P, Fornari L, Caruso A, Varone G, Genazzani AR. Genomic and non-genomic effects of estrogens on endothelial cells. *Steroids*. 2004;69(8):537-42.
203. Menazza S, Murphy E. The Expanding Complexity of Estrogen Receptor Signaling in the Cardiovascular System. *Circulation Research*. 2016;118(6):994-1007.
204. Acconcia F, Totta P, Ogawa S, Cardillo I, Inoue S, Leone S, et al. Survival versus apoptotic 17 $\beta$ -estradiol effect: Role of ER $\alpha$  and ER $\beta$  activated non-genomic signaling. *Journal of Cellular Physiology*. 2005;203(1):193-201.
205. Chen JQ, Delannoy M, Cooke C, Yager JD. Mitochondrial localization of ER $\alpha$  and ER $\beta$  in human MCF7 cells. *American Journal of Physiology-Endocrinology and Metabolism*. 2004;286(6):E1011-E22.
206. Chambliss KL, Yuhanna IS, Mineo C, Liu P, German Z, Sherman TS, et al. Estrogen receptor  $\alpha$  and endothelial nitric oxide synthase are organized into a functional signaling module in caveolae. *Circ Res*. 2000;87(11):E44-52.
207. Bulger M, Groudine M. Functional and Mechanistic Diversity of Distal Transcription Enhancers. *Cell*. 2011;144(3):327-39.

208. Rao Suhas SP, Huntley Miriam H, Durand Neva C, Stamenova Elena K, Bochkov Ivan D, Robinson James T, et al. A 3D Map of the Human Genome at Kilobase Resolution Reveals Principles of Chromatin Looping. *Cell*. 2014;159(7):1665-80.
209. Fullwood MJ, Liu MH, Pan YF, Liu J, Xu H, Mohamed YB, et al. An oestrogen-receptor-alpha-bound human chromatin interactome. *Nature*. 2009;462(7269):58-64.
210. Pan YF, Wansa KDSA, Liu MH, Zhao B, Hong SZ, Tan PY, et al. Regulation of Estrogen Receptor-mediated Long Range Transcription via Evolutionarily Conserved Distal Response Elements\*. *Journal of Biological Chemistry*. 2008;283(47):32977-88.
211. Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, et al. Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature*. 2012;485(7398):376-80.
212. Mourad R, Hsu P-Y, Juan L, Shen C, Koneru P, Lin H, et al. Estrogen Induces Global Reorganization of Chromatin Structure in Human Breast Cancer Cells. *PLOS ONE*. 2014;9(12):e113354.
213. Rada-Iglesias A, Bajpai R, Swigut T, Brugmann SA, Flynn RA, Wysocka J. A unique chromatin signature uncovers early developmental enhancers in humans. *Nature*. 2011;470(7333):279-83.
214. Bonn S, Zinzen RP, Girardot C, Gustafson EH, Perez-Gonzalez A, Delhomme N, et al. Tissue-specific analysis of chromatin state identifies temporal signatures of enhancer activity during embryonic development. *Nature genetics*. 2012;44(2):148-56.
215. Lovén J, Hoke Heather A, Lin Charles Y, Lau A, Orlando David A, Vakoc Christopher R, et al. Selective Inhibition of Tumor Oncogenes by Disruption of Super-Enhancers. *Cell*. 2013;153(2):320-34.
216. Pott S, Lieb JD. What are super-enhancers? *Nature genetics*. 2015;47(1):8-12.
217. Field A, Adelman K. Evaluating Enhancer Function and Transcription. *Annual Review of Biochemistry*. 2020;89(1):213-34.
218. Wang D, Garcia-Bassets I, Benner C, Li W, Su X, Zhou Y, et al. Reprogramming transcription by distinct classes of enhancers functionally defined by eRNA. *Nature*. 2011;474(7351):390-4.
219. Andersson R, Gebhard C, Miguel-Escalada I, Hoof I, Bornholdt J, Boyd M, et al. An atlas of active enhancers across human cell types and tissues. *Nature*. 2014;507(7493):455-61.
220. Lai F, Orom UA, Cesaroni M, Beringer M, Taatjes DJ, Blobel GA, et al. Activating RNAs associate with Mediator to enhance chromatin architecture and transcription. *Nature*. 2013;494(7438):497-501.
221. Pennacchio LA, Bickmore W, Dean A, Nobrega MA, Bejerano G. Enhancers: five essential questions. *Nature reviews Genetics*. 2013;14(4):288-95.
222. RNA-seq of 934 human cancer cell lines from the Cancer Cell Line Encyclopedia, arrayexpress-repository, V1. <https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-2770>. (2013).
223. Baker M. Reproducibility: Seek out stronger science. *Nature*. 2016;537(7622):703-4.
224. Taussig MJ, Fonseca C, Trimmer JS. Antibody validation: a view from the mountains. *N Biotechnol*. 2018;45:1-8.

225. Andersson S, Sundberg M, Pristovsek N, Ibrahim A, Jonsson P, Katona B, et al. Insufficient antibody validation challenges oestrogen receptor beta research. *Nature communications*. 2017;8:15840.
226. Hases L, Indukuri R, Birgersson M, Nguyen-Vu T, Lozano R, Saxena A, et al. Intestinal estrogen receptor beta suppresses colon inflammation and tumorigenesis in both sexes. *Cancer Letters*. 2020.
227. Nelson AW, Groen AJ, Miller JL, Warren AY, Holmes KA, Tarulli GA, et al. Comprehensive assessment of estrogen receptor beta antibodies in cancer cell line models and tissue reveals critical limitations in reagent specificity. *Molecular and cellular endocrinology*. 2017;440:138-50.
228. Nelson RL, Dollear T, Freels S, Persky V. The relation of age, race, and gender to the subsite location of colorectal carcinoma. *Cancer*. 1997;80(2):193.
229. Hendifar A, Yang D, Lenz F, Lurje G, Pohl A, Lenz C, et al. Gender Disparities in Metastatic Colorectal Cancer Survival. *Clinical Cancer Research*. 2009;15(20):6391-7.
230. Wichmann MW, Müller C, Hornung HM, Lau-Werner U, Schildberg FW. Gender differences in long-term survival of patients with colorectal cancer. *British Journal of Surgery*. 2001;88(8):1092-8.
231. McArdle CS, McMillan DC, Hole DJ. Male gender adversely affects survival following surgery for colorectal cancer. *BJS*. 2003;90(6):711-5.
232. Chlebowski RT, Wactawski-Wende J, Ritenbaugh C, Hubbell FA, Ascensao J, Rodabough RJ, et al. Estrogen plus Progestin and Colorectal Cancer in Postmenopausal Women. *New England Journal of Medicine*. 2004;350(10):991-1004.
233. Caiazza F, Ryan EJ, Doherty G, Winter DC, Sheahan K. Estrogen Receptors and Their Implications in Colorectal Carcinogenesis. *Frontiers in Oncology*. 2015;5(19).
234. Simin J, Tamimi R, Lagergren J, Adami H-O, Brusselsaers N. Menopausal hormone therapy and cancer risk: An overestimated risk? *European Journal of Cancer*. 2017;84:60-8.
235. Botteri E, Stør NC, Sakshaug S, Graff-Iversen S, Vangen S, Hofvind S, et al. Menopausal hormone therapy and colorectal cancer: a linkage between nationwide registries in Norway. *BMJ open*. 2017;7(11):e017639-e.
236. Luan NN, Wu L, Gong TT, Wang YL, Lin B, Wu QJ. Nonlinear reduction in risk for colorectal cancer by oral contraceptive use: a meta-analysis of epidemiological studies. *Cancer Causes Control*. 2015;26(1):65-78.
237. Ganfeng Luo, Yanting Zhang, Li Wang, Yuanwei Huang, Qiuyan Yu, Pi Guo, et al. Risk of colorectal cancer with hysterectomy and oophorectomy: A systematic review and meta-analysis. *International Journal of Surgery*, 2016(34):88-95.
238. Yan L, Spitznagel EL, Bosland MC. Soy consumption and colorectal cancer risk in humans: a meta-analysis. *Cancer Epidemiol Biomarkers Prev*. 2010;19(1):148-58.
239. Flood DM, Weiss NS, Cook LS, Emerson JC, Schwartz SM, Potter JD. Colorectal cancer incidence in Asian migrants to the United States and their descendants. *Cancer Causes Control*. 2000;11(5):403-11.
240. Kim S-E, Shim K-N, Jung S-A, Yoo K, Moon IH. An association between obesity and the prevalence of colonic adenoma according to age and gender. *Journal of Gastroenterology*. 2007;42(8):616-23.

241. Williams C, DiLeo A, Niv Y, Gustafsson J-Å. Estrogen receptor beta as target for colorectal cancer prevention. *Cancer Letters*. 2016;372(1):48-56.
242. Guo JY, Li X, Browning JD, Jr., Rottinghaus GE, Lubahn DB, Constantinou A, et al. Dietary soy isoflavones and estrone protect ovariectomized ERalphaKO and wild-type mice from carcinogen-induced colon cancer. *The Journal of nutrition*. 2004;134(1):179-82.
243. Javid SH, Moran AE, Carothers AM, Redston M, Bertagnolli MM. Modulation of tumor formation and intestinal cell migration by estrogens in the Apc(Min/+) mouse model of colorectal cancer. *Carcinogenesis*. 2005;26(3):587-95.
244. Weige CC, Allred KF, Allred CD. Estradiol alters cell growth in nonmalignant colonocytes and reduces the formation of preneoplastic lesions in the colon. *Cancer Res*. 2009;69(23):9118-24.
245. Nguyen-Vu T, Wang J, Mesmar F, Mukhopadhyay S, Saxena A, McCollum CW, et al. Estrogen receptor beta reduces colon cancer metastasis through a novel miR-205 - PROX1 mechanism. *Oncotarget*. 2016;7(27):42159.
246. Saleiro D, Murillo G, Benya RV, Bissonnette M, Hart J, Mehta RG. Estrogen receptor- $\beta$  protects against colitis-associated neoplasia in mice. *International journal of cancer*. 2012;131(11):2553-61.
247. Heijmans J, Wielenga MC, Rosekrans SL, van Lidth de Jeude JF, Roelofs J, Groothuis P, et al. Oestrogens promote tumorigenesis in a mouse model for colitis-associated cancer. *Gut*. 2014;63(2):310-6.
248. Armstrong CM, Allred KF, Weeks BR, Chapkin RS, Allred CD. Estradiol Has Differential Effects on Acute Colonic Inflammation in the Presence and Absence of Estrogen Receptor  $\beta$  Expression. *Digestive Diseases and Sciences*. 2017;62(8):1977-84.
249. Giroux V, Lemay F, Bernatchez G, Robitaille Y, Carrier JC. Estrogen receptor  $\beta$  deficiency enhances small intestinal tumorigenesis in ApcMin/+ mice. *International journal of cancer*. 2008;123(2):303-11.
250. Hartman J, Edvardsson K, Lindberg K, Zhao C, Williams C, Ström A, et al. Tumor Repressive Functions of Estrogen Receptor  $\beta$  in SW480 Colon Cancer Cells. *Cancer Research*. 2009;69(15):6100-6.
251. Edvardsson K, Strom A, Jonsson P, Gustafsson J-Å, Williams C. Estrogen Receptor  $\beta$  Induces Antiinflammatory and Antitumorigenic Networks in Colon Cancer Cells 2011. 969-79 p.
252. Hsu HH, Cheng SF, Wu CC, Chu CH, Weng YJ, Lin CS, et al. Apoptotic effects of over-expressed estrogen receptor-beta on LoVo colon cancer cell is mediated by p53 signalings in a ligand-dependent manner. *Chin J Physiol*. 2006;49(2):110-6.
253. Heldring N, Pike A, Andersson S, Matthews J, Cheng G, Hartman J, et al. Estrogen receptors: how do they signal and what are their targets. *Physiol Rev*. 2007;87(3):905-31.
254. Loven MA, Likhite VS, Choi I, Nardulli AM. Estrogen response elements alter coactivator recruitment through allosteric modulation of estrogen receptor beta conformation. *The Journal of biological chemistry*. 2001;276(48):45282-8.
255. Loven MA, Wood JR, Nardulli AM. Interaction of estrogen receptors alpha and beta with estrogen response elements. *Molecular and cellular endocrinology*. 2001;181(1-2):151-63.
256. Cowley SM, Parker MG. A comparison of transcriptional activation by ER alpha and ER beta. *The Journal of steroid biochemistry and molecular biology*. 1999;69(1-6):165.



257. Deroo BJ, Buensuceso AV. Minireview: Estrogen receptor-beta: mechanistic insights from recent studies. *Molecular endocrinology* (Baltimore, Md). 2010;24(9):1703-14.
258. Iwafuchi-Doi M, Donahue G, Kakumanu A, Watts Jason A, Mahony S, Pugh BF, et al. The Pioneer Transcription Factor FoxA Maintains an Accessible Nucleosome Configuration at Enhancers for Tissue-Specific Gene Activation. *Molecular cell*. 2016;62(1):79-91.
259. Hua H, Zhang H, Kong Q, Jiang Y. Mechanisms for estrogen receptor expression in human cancer. *Experimental Hematology & Oncology*. 2018;7(1):24.
260. Zhou Y, Zeng C, Li X, Wu PL, Yin L, Yu XL, et al. IGF-I stimulates ER $\beta$  and aromatase expression via IGF1R/PI3K/AKT-mediated transcriptional activation in endometriosis. *J Mol Med (Berl)*. 2016;94(8):887-97.
261. Zhang X, Leung YK, Ho SM. AP-2 regulates the transcription of estrogen receptor (ER)-beta by acting through a methylation hotspot of the 0N promoter in prostate cancer cells. *Oncogene*. 2007;26(52):7346-54.
262. Cai W, Rambaud J, Teboul M, Masse I, Benoit G, Gustafsson J-Å, et al. Expression Levels of Estrogen Receptor  $\beta$  Are Modulated by Components of the Molecular Clock. *Molecular and Cellular Biology*. 2008;28(2):784-93.
263. Wang L, Ozark PA, Smith ER, Zhao Z, Marshall SA, Rendleman EJ, et al. TET2 coactivates gene expression through demethylation of enhancers. *Sci Adv*. 2018;4(11):eaau6986-eaau.
264. Broome R, Chernukhin I, Jamieson S, Kishore K, Papachristou EK, Mao SQ, et al. TET2 is a component of the estrogen receptor complex and controls 5mC to 5hmC conversion at estrogen receptor cis-regulatory regions. *Cell reports*. 2021;34(8):108776.
265. Cerciati M, Unkila M, Garcia-Segura LM, Arevalo MA. Selective estrogen receptor modulators decrease the production of interleukin-6 and interferon-gamma-inducible protein-10 by astrocytes exposed to inflammatory challenge in vitro. *Glia*. 2010;58(1):93-102.
266. Cvaro A, Tzagarakis-Foster C, Tatomer D, Paruthiyil S, Fox MS, Leitman DC. Distinct roles of unliganded and liganded estrogen receptors in transcriptional repression. *Molecular cell*. 2006;21(4):555-64.
267. Kalaitzidis D, Gilmore TD. Transcription factor cross-talk: the estrogen receptor and NF-kappaB. *Trends in endocrinology and metabolism: TEM*. 2005;16(2):46-52.
268. Nettles KW, Gil G, Nowak J, Metivier R, Sharma VB, Greene GL. CBP Is a dosage-dependent regulator of nuclear factor-kappaB suppression by the estrogen receptor. *Molecular endocrinology* (Baltimore, Md). 2008;22(2):263-72.
269. Adamson AD, Friedrichsen S, Semprini S, Harper CV, Mullins JJ, White MR, et al. Human prolactin gene promoter regulation by estrogen: convergence with tumor necrosis factor-alpha signaling. *Endocrinology*. 2008;149(2):687-94.
270. King AE, Collins F, Klonisch T, Sallenave JM, Critchley HO, Saunders PT. An additive interaction between the NFkappaB and estrogen receptor signalling pathways in human endometrial epithelial cells. *Human reproduction* (Oxford, England). 2010;25(2):510-8.
271. Rubio MF, Werbajh S, Cafferata EG, Quaglini A, Colo GP, Nojek IM, et al. TNF-alpha enhances estrogen-induced cell proliferation of estrogen-dependent breast tumor cells through a complex containing nuclear factor-kappa B. *Oncogene*. 2006;25(9):1367-77.

272. Wissink S, van der Burg B, Katzenellenbogen BS, van der Saag PT. Synergistic activation of the serotonin-1A receptor by nuclear factor-kappa B and estrogen. *Molecular endocrinology* (Baltimore, Md). 2001;15(4):543-52.
273. Stender JD, Nwachukwu JC, Kastrati I, Kim Y, Strid T, Yakir M, et al. Structural and Molecular Mechanisms of Cytokine-Mediated Endocrine Resistance in Human Breast Cancer Cells. *Molecular cell*. 2017;65(6):1122-35.e5.
274. Pradhan M, Baumgarten SC, Bembinster LA, Frasor J. CBP Mediates NF- $\kappa$ B-Dependent Histone Acetylation and Estrogen Receptor Recruitment to an Estrogen Response Element in the BIRC3 Promoter. *Molecular and Cellular Biology*. 2012;32(2):569-75.
275. Franco HL, Nagari A, Kraus WL. TNF $\alpha$  signaling exposes latent estrogen receptor binding sites to alter the breast cancer cell transcriptome. *Molecular cell*. 2015;58(1):21-34.
276. Hah N, Murakami S, Nagari A, Danko CG, Kraus WL. Enhancer transcripts mark active estrogen receptor binding sites. *Genome research*. 2013;23(8):1210-23.
277. Pradhan M, Baumgarten SC, Bembinster LA, Frasor J. CBP mediates NF- $\kappa$ B-dependent histone acetylation and estrogen receptor recruitment to an estrogen response element in the BIRC3 promoter. *Molecular and cellular biology*. 2012;32(2):569-75.
278. Stender JD, Nwachukwu JC, Kastrati I, Kim Y, Strid T, Yakir M, et al. Structural and Molecular Mechanisms of Cytokine-Mediated Endocrine Resistance in Human Breast Cancer Cells. *Molecular cell*. 2017;65(6):1122-35.e5.
279. Edvardsson K, Strom A, Jonsson P, Gustafsson JA, Williams C. Estrogen receptor beta induces antiinflammatory and antitumorigenic networks in colon cancer cells. *Molecular endocrinology* (Baltimore, Md). 2011;25(6):969-79.
280. Flach KD, Zwart W. The first decade of estrogen receptor cistromics in breast cancer. *Journal of Endocrinology*. 2016;229(2):R43-R56.
281. Welboren W-J, van Driel MA, Janssen-Megens EM, van Heeringen SJ, Sweep FCGJ, Span PN, et al. ChIP-Seq of ER $\alpha$  and RNA polymerase II defines genes differentially responding to ligands. *The EMBO Journal*. 2009;28(10):1418-28.
282. Lupien M, Eeckhoutte J, Meyer CA, Wang Q, Zhang Y, Li W, et al. FoxA1 translates epigenetic signatures into enhancer-driven lineage-specific transcription. *Cell*. 2008;132:958-70.
283. Liu Y, Gao H, Marstrand TT, Strom A, Valen E, Sandelin A, et al. The genome landscape of ER $\alpha$ - and ER $\beta$ -binding DNA regions. *Proceedings of the National Academy of Sciences of the United States of America*. 2008;105(7):2604-9.
284. Grober OMV, Mutarelli M, Giurato G, Ravo M, Cicatiello L, De Filippo MR, et al. Global analysis of estrogen receptor beta binding to breast cancer cell genome reveals an extensive interplay with estrogen receptor alpha for target gene regulation. *BMC Genomics*. 2011;12:36-.
285. Madak-Erdogan Z, Charn T-H, Jiang Y, Liu ET, Katzenellenbogen JA, Katzenellenbogen BS. Integrative genomics of gene and metabolic regulation by estrogen receptors  $\alpha$  and  $\beta$ , and their coregulators. *Mol Syst Biol*. 2013;9:676-.
286. Vivar OI, Zhao X, Saunier EF, Griffin C, Mayba OS, Tagliaferri M, et al. Estrogen receptor beta binds to and regulates three distinct classes of target genes. *The Journal of biological chemistry*. 2010;285(29):22059-66.

287. Raut S, Kumar AV, Khambata K, Deshpande S, Balasinor NH. Genome-wide identification of estrogen receptor binding sites reveals novel estrogen-responsive pathways in adult male germ cells. *Biochem J.* 2020;477(12):2115-31.
288. Han SJ, Lee JE, Cho YJ, Park MJ, O'Malley BW. Genomic Function of Estrogen Receptor  $\beta$  in Endometriosis. *Endocrinology.* 2019;160(11):2495-516.
289. Karnuta JM, Scacheri PC. Enhancers: bridging the gap between gene control and human disease. *Human Molecular Genetics.* 2018;27(R2):R219-R27.
290. Zurlo J, Rudacille D, Goldberg AM. The three Rs: the way forward. *Environ Health Perspect.* 1996;104(8):878-80.
291. Rumi MAK, Singh P, Roby KF, Zhao X, Iqbal K, Ratri A, et al. Defining the Role of Estrogen Receptor  $\beta$  in the Regulation of Female Fertility. *Endocrinology.* 2017;158(7):2330-43.
292. De Robertis M, Massi E, Poeta ML, Carotti S, Morini S, Cecchetelli L, et al. The AOM/DSS murine model for the study of colon carcinogenesis: From pathways to diagnosis and therapy studies. *J Carcinog.* 2011;10:9.
293. Snider AJ, Bialkowska AB, Ghaleb AM, Yang VW, Obeid LM, Hannun YA. Murine Model for Colitis-Associated Cancer of the Colon. *Methods in molecular biology (Clifton, NJ).* 2016;1438:245-54.
294. De Robertis M, Massi E, Poeta ML, Carotti S, Morini S, Cecchetelli L, et al. The AOM/DSS murine model for the study of colon carcinogenesis: From pathways to diagnosis and therapy studies. *Journal of carcinogenesis.* 2011;10:9-.
295. Karunanithi S, Levi L. High-fat diet and colorectal cancer: myths and facts. *Future Oncology.* 2018;14(6):493-5.
296. Martínez-Maqueda D, Miralles B, Recio I. HT29 Cell Line. In: Verhoeckx K, Cotter P, López-Expósito I, Kleiveland C, Lea T, Mackie A, et al., editors. *The Impact of Food Bioactives on Health: in vitro and ex vivo models.* Cham: Springer International Publishing; 2015. p. 113-24.
297. Leibovitz A, Stinson JC, McCombs WB, 3rd, McCoy CE, Mazur KC, Mabry ND. Classification of human colorectal adenocarcinoma cell lines. *Cancer Res.* 1976;36(12):4562-9.
298. Ahmed D, Eide PW, Eilertsen IA, Danielsen SA, Eknæs M, Hektoen M, et al. Epigenetic and genetic features of 24 colon cancer cell lines. *Oncogenesis.* 2013;2(9):e71-e.
299. Edvardsson K, Nguyen-Vu T, Kalasekar SM, Ponten F, Gustafsson JA, Williams C. Estrogen receptor beta expression induces changes in the microRNA pool in human colon cancer cells. *Carcinogenesis.* 2013;34(7):1431-41.
300. Nakato R, Shirahige K. Recent advances in ChIP-seq analysis: from quality management to whole-genome annotation. *Briefings in bioinformatics.* 2017;18(2):279-90.
301. Furey TS. ChIP-seq and beyond: new and improved methodologies to detect and characterize protein-DNA interactions. *Nature Reviews Genetics.* 2012;13:840.
302. Park PJ. ChIP-seq: advantages and challenges of a maturing technology. *Nature Reviews Genetics.* 2009;10:669.
303. Jiang S, Mortazavi A. Integrating ChIP-seq with other functional genomics data. *Briefings in functional genomics.* 2018;17(2):104-15.

304. Lin TH, Pajarinen J, Lu L, Nabeshima A, Cordova LA, Yao Z, et al. NF- $\kappa$ B as a Therapeutic Target in Inflammatory-Associated Bone Diseases. *Advances in protein chemistry and structural biology*. 2017;107:117-54.
305. Sheaffer KL, Schug J. ChIP-Seq: Library Preparation and Sequencing. *Methods in molecular biology* (Clifton, NJ). 2016;1402:101-17.
306. Rehim R, Bartusel M, Solinas F, Altmüller J, Rada-Iglesias A. Chromatin Immunoprecipitation (ChIP) Protocol for Low-abundance Embryonic Samples. *Journal of visualized experiments : JoVE*. 2017(126).
307. Beischlag TV, Prefontaine GG, Hankinson O. ChIP-re-ChIP: Co-occupancy Analysis by Sequential Chromatin Immunoprecipitation. *Methods in molecular biology* (Clifton, NJ). 2018;1689:103-12.
308. Ribarska T, Gilfillan GD. Native Chromatin Immunoprecipitation-Sequencing (ChIP-Seq) from Low Cell Numbers. *Methods in molecular biology* (Clifton, NJ). 2018;1689:157-66.
309. Diaz RE, Sanchez A, Anton Le Berre V, Bouet JY. High-Resolution Chromatin Immunoprecipitation: ChIP-Sequencing. *Methods in molecular biology* (Clifton, NJ). 2017;1624:61-73.
310. Chen Y, Negre N, Li Q, Mieczkowska JO, Slattery M, Liu T, et al. Systematic evaluation of factors influencing ChIP-seq fidelity. *Nature methods*. 2012;9(6):609-14.
311. Khrameeva EE, Gelfand MS. Biases in read coverage demonstrated by interlaboratory and interplatform comparison of 117 mRNA and genome sequencing experiments. *BMC bioinformatics*. 2012;13 Suppl 6:S4.
312. Minoche AE, Dohm JC, Himmelbauer H. Evaluation of genomic high-throughput sequencing data generated on Illumina HiSeq and genome analyzer systems. *Genome biology*. 2011;12(11):R112.
313. Schwartz S, Oren R, Ast G. Detection and removal of biases in the analysis of next-generation sequencing reads. *PLoS One*. 2011;6(1):e16685.
314. Orlando DA, Chen MW, Brown VE, Solanki S, Choi YJ, Olson ER, et al. Quantitative ChIP-Seq normalization reveals global modulation of the epigenome. *Cell reports*. 2014;9(3):1163-70.
315. Bonhoure N, Bounova G, Bernasconi D, Praz V, Lammers F, Canella D, et al. Quantifying ChIP-seq data: a spiking method providing an internal reference for sample-to-sample normalization. *Genome research*. 2014;24(7):1157-68.
316. Guertin MJ, Cullen AE, Markowitz F, Holding AN. Parallel factor ChIP provides essential internal control for quantitative differential ChIP-seq. *Nucleic acids research*. 2018;46(12):e75-e.
317. Chen TW, Li HP, Lee CC, Gan RC, Huang PJ, Wu TH, et al. ChIPseek, a web-based analysis tool for ChIP data. *BMC Genomics*. 2014;15(1):539.
318. Boeva V, Lermine A, Barette C, Guillouf C, Barillot E. Nebula--a web-server for advanced ChIP-seq data analysis. *Bioinformatics*. 2012;28(19):2517-9.
319. Li M, Tang L, Wu F-X, Pan Y, Wang J. CSA: a web service for the complete process of ChIP-Seq analysis. *BMC Bioinformatics*. 2019;20(15):515.

320. Rotem A, Ram O, Shores N, Sperling RA, Goren A, Weitz DA, et al. Single-cell ChIP-seq reveals cell subpopulations defined by chromatin state. *Nature biotechnology*. 2015;33(11):1165-72.
321. Grosselin K, Durand A, Marsolier J, Poitou A, Marangoni E, Nemati F, et al. High-throughput single-cell ChIP-seq identifies heterogeneity of chromatin states in breast cancer. *Nature genetics*. 2019;51(6):1060-6.
322. Giroux V, Bernatchez G, Carrier JC. Chemopreventive effect of ERbeta-Selective agonist on intestinal tumorigenesis in Apc(Min/+) mice. *Mol Carcinog*. 2011;50(5):359-69.
323. Giroux V, Lemay F, Bernatchez G, Robitaille Y, Carrier JC. Estrogen receptor beta deficiency enhances small intestinal tumorigenesis in ApcMin/+ mice. *International journal of cancer*. 2008;123(2):303-11.
324. Saleiro D, Murillo G, Lubahn DB, Kopelovich L, Korach KS, Mehta RG. Enhanced induction of mucin-depleted foci in estrogen receptor {beta} knockout mice. *Cancer prevention research (Philadelphia, Pa)*. 2010;3(9):1198-204.
325. Saleiro D, Murillo G, Benya RV, Bissonnette M, Hart J, Mehta RG. Estrogen receptor-beta protects against colitis-associated neoplasia in mice. *International journal of cancer*. 2012;131(11):2553-61.
326. Martinetti V, Picariello L, Tognarini I, Carbonell Sala S, Gozzini A, Azzari C, et al. ERbeta is a potent inhibitor of cell proliferation in the HCT8 human colon cancer cell line through regulation of cell cycle components. *Endocr Relat Cancer*. 2005;12(2):455-69.
327. Hartman J, Edvardsson K, Lindberg K, Zhao C, Williams C, Strom A, et al. Tumor repressive functions of estrogen receptor beta in SW480 colon cancer cells. *Cancer Res*. 2009;69(15):6100-6.
328. Bielecki A, Roberts J, Mehta R, Raju J. Estrogen receptor-beta mediates the inhibition of DLD-1 human colon adenocarcinoma cells by soy isoflavones. *Nutr Cancer*. 2011;63(1):139-50.
329. Tu Z, Ma Y, Tian J, Li H, Akers W, Achilefu S, et al. Estrogen receptor beta potentiates the antiproliferative effect of raloxifene and affects the cell migration and invasion in HCT-116 colon cancer cells. *Journal of cancer research and clinical oncology*. 2012;138(7):1091-103.
330. Ibrahim A, Hugerth LW, Hases L, Saxena A, Seifert M, Thomas Q, et al. Colitis-induced colorectal cancer and intestinal epithelial estrogen receptor beta impact gut microbiota diversity. *International journal of cancer*. 2018.
331. Doumpas N, Lampart F, Robinson MD, Lentini A, Nestor CE, Cantù C, et al. TCF/LEF dependent and independent transcriptional regulation of Wnt/ $\beta$ -catenin target genes. *Embo j*. 2019;38(2).
332. Becker WR, Nevins SA, Chen DC, Chiu R, Horning A, Laquindanum R, et al. Single-cell analyses reveal a continuum of cell state and composition changes in the malignant transformation of polyps to colorectal cancer. *bioRxiv*. 2021:2021.03.24.436532.
333. Wei D, Kanai M, Huang S, Xie K. Emerging role of KLF4 in human gastrointestinal cancer. *Carcinogenesis*. 2006;27(1):23-31.
334. Indukuri R, Jafferli MH, Song D, Damdimopoulos A, Hases L, Zhao C, et al. Genome-wide estrogen receptor  $\beta$  chromatin binding in human colon cancer cells reveals its tumor suppressor activity. *International journal of cancer*. 2021:1-15.

335. Cox AJ, West NP, Cripps AW. Obesity, inflammation, and the gut microbiota. *Lancet Diabetes Endocrinol.* 2015;3(3):207-15.
336. Chen J, Iverson D. Estrogen in obesity-associated colon cancer: friend or foe? Protecting postmenopausal women but promoting late-stage colon cancer. *Cancer Causes & Control.* 2012;23(11):1767-73.
337. Savva C, Korach-André M. Estrogen Receptor beta (ER $\beta$ ) Regulation of Lipid Homeostasis-Does Sex Matter? *Metabolites.* 2020;10(3):116.
338. Giatromanolaki A, Sivridis E, Maltezos E, Papazoglou D, Simopoulos C, Gatter KC, et al. Hypoxia inducible factor 1 $\alpha$  and 2 $\alpha$  overexpression in inflammatory bowel disease. *Journal of Clinical Pathology.* 2003;56(3):209-13.
339. Geary N, Asarian L, Korach KS, Pfaff DW, Ogawa S. Deficits in E2-dependent control of feeding, weight gain, and cholecystokinin satiation in ER- $\alpha$  null mice. *Endocrinology.* 2001;142(11):4751-7.
340. Heine PA, Taylor JA, Iwamoto GA, Lubahn DB, Cooke PS. Increased adipose tissue in male and female estrogen receptor- $\alpha$  knockout mice. *Proceedings of the National Academy of Sciences.* 2000;97(23):12729-34.
341. Hases L, Archer A, Indukuri R, Birgersson M, Savva C, Korach-André M, et al. High-fat diet and estrogen impacts the colon and its transcriptome in a sex-dependent manner. *Scientific Reports.* 2020;10(1):16160.
342. Jeong S-J, Radonovich M, Brady JN, Pise-Masison CA. HTLV-I Tax induces a novel interaction between p65/RelA and p53 that results in inhibition of p53 transcriptional activity. *Blood.* 2004;104(5):1490-7.
343. Schneider G, Henrich A, Greiner G, Wolf V, Lovas A, Wieczorek M, et al. Cross talk between stimulated NF- $\kappa$ B and the tumor suppressor p53. *Oncogene.* 2010;29(19):2795-806.
344. Frank AK, Leu JI-J, Zhou Y, Devarajan K, Nedelko T, Klein-Szanto A, et al. The Codon 72 Polymorphism of p53 Regulates Interaction with NF- $\kappa$ B and Transactivation of Genes Involved in Immunity and Inflammation. *Molecular and Cellular Biology.* 2011;31(6):1201-13.
345. Rahnamoun H, Lu H, Duttke SH, Benner C, Glass CK, Lauberth SM. Mutant p53 shapes the enhancer landscape of cancer cells in response to chronic immune signaling. *Nature communications.* 2017;8(1):754.
346. Boynton FDD, Ericsson AC, Uchihashi M, Dunbar ML, Wilkinson JE. Doxycycline induces dysbiosis in female C57BL/6NCrl mice. *BMC research notes.* 2017;10(1):644-.
347. Hou X, Zhu L, Zhang X, Zhang L, Bao H, Tang M, et al. Testosterone disruptor effect and gut microbiome perturbation in mice: Early life exposure to doxycycline. *Chemosphere.* 2019;222:722-31.
348. Mohammed H, Taylor C, Brown GD, Papachristou EK, Carroll JS, D'Santos CS. Rapid immunoprecipitation mass spectrometry of endogenous proteins (RIME) for analysis of chromatin complexes. *Nat Protoc.* 2016;11(2):316-26.
349. Papachristou EK, Kishore K, Holding AN, Harvey K, Roumeliotis TI, Chilamakuri CSR, et al. A quantitative mass spectrometry-based approach to monitor the dynamics of endogenous chromatin-associated protein complexes. *Nature communications.* 2018;9(1):2311.

350. Tremblay A, Tremblay GB, Labrie F, Giguère V. Ligand-Independent Recruitment of SRC-1 to Estrogen Receptor  $\beta$  through Phosphorylation of Activation Function AF-1. *Molecular cell*. 1999;3(4):513-9.
351. Tremblay A, Giguère V. Contribution of steroid receptor coactivator-1 and CREB binding protein in ligand-independent activity of estrogen receptor beta. *The Journal of steroid biochemistry and molecular biology*. 2001;77(1):19-27.
352. Hases L, Ibrahim A, Chen X, Liu Y, Hartman J, Williams C. The Importance of Sex in the Discovery of Colorectal Cancer Prognostic Biomarkers. *International journal of molecular sciences*. 2021;22(3).